INSTRUCTIONAL CHECKLIST:



Bedside Mix – Eclipse Ball Pre-Infusion Preparation

https://www.youtube.com/watch?v=rJI8xurPJdI&feature=emb logo

For additional education please contact your agency for assistance

Four Steps for Performing Infusions:

roal steps for refronting infusions.			
Pra	actice good hand-washing 1 Prepare supplies 2 Draw up the medication into 4		
Med	lication(s) to infuse: Home Infusion Nurse (name, #):		
	Home Infusion Pharmacist (name, #):		
	ice good Always wash hands Soap and water Hand sanitizer Wash for a full 20 -washing when instructed preferred (optional) seconds or rub until dry		
	STEP 1: PRACTICE GOOD HAND-WASHING		
	 Soap and water is always preferred. If soap and water are not available, hand sanitizer is acceptable. Make it a habit to wet your hands, apply soap, and lather for 20 seconds (hum the Happy Birthday song twice). If using hand sanitizer, rub a dime-sized amount between your hands until dry. Never wave or blow on your hands to speed up drying. Remember, if you are using gloves, you must still wash your hands. Washing your hands helps reduce your chance of infection. 		
Pre	Page Supplies Note infusion Start time WASH HANDS WASH Start time WASH WASH Supplies WASH HANDS WASH MANDS Supplies		
	STEP 2: PREPARE SUPPLIES		
	Note the time to start the infusion: AM PM		
	WASH YOUR HANDS.		
	 Soap and water are preferred, hand sanitizer optional if soap and water are not available. Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry. 		
	Clean your workspace with cleaning solution (alcohol/sanitizing wipes or Lysol + paper towels).		



	Select a dry, hard work surface to set up your medication and supplies.	
	Gather the vials of medication and mixing fluid and place them in your work area. You will need your orders as well.	
	 Double-check your medication to confirm the following information before the infusion: Medication has my name and date of birth. Medication name is correct. Medication dose is correct. Medication is not expired. 	
	 WASH YOUR HANDS. ❖ Soap and water are preferred, hand sanitizer optional if soap and water are not available. ❖ Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry. 	
	Gather the following supplies and place on your workspace (these are for mixing your medications):	
Draw up the medication WASH HANDS Prepare vials Prepare syringe and needle Withdraw diluent Mix medication Withdraw final medication		
	STEP 3: DRAW UP THE MEDICATION	
	 WASH YOUR HANDS. ❖ Soap and water are preferred, hand sanitizer optional if soap and water are not available. ❖ Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry. 	
	Remove the caps from each vial of medication and mixing fluid. Remember the number of vials and mixing fluid needed will be dependent upon the dose of infusion and dose in each vial. I will need vials of medication and vials of mixing fluid. Wipe the top of each vial with 1 alcohol pad per vial.	
	Prepare your syringe and needle. ❖ Open the package and remove the syringe and needle. ❖ If the needle is not attached to the syringe, attach the needle to the syringe by screwing the needle to the end of the syringe careful not to touch the tip of the syringe or bottom of the needle. ❖ Pull back on the plunger of the syringe filling it with air to the volume of diluent you intend to withdraw.	
	 Withdraw mixing fluid. Remove the plastic cover over the needle and place on the work area within reach. Insert the needle into the top of the mix vial. Pick up the bottle of mixing fluid with one hand while holding the syringe in place with the other and invert the vial upside down. 	



 Slowly inject air into the vial with slight pressure and allow the pressure in the vial to push fluid back into the syringe as the plunger retracts. Remember to keep the tip of the needle submerged in the diluent. Continue to draw up mixing fluid until all the fluid is withdrawn into the syringe. Remove the syringe from the mix vial and set vial aside.
Mix the medication by Inserting the needle into the medication vial and slowly injecting the diluent into the medication vial. Leave the needle in the vial while you gently swirl the vial. Do not aggressively shake the vial.
 Withdraw medication. When you are certain all the powder medication within the vial has dissolved, invert the vial of medication. Pull back on the syringe to withdraw the required volume of medication. You may need to adjust the needle's location within the vial to withdraw all of the medication included. Once you have withdrawn all the medication, remove the syringe from the vial. Carefully unscrew the needle from the syringe and place in a sharps container. Sit the syringe aside careful not to contaminate the end of the syringe.
 ct medication Locate the Attach Inject syringe Close the device cap syringe contents device HANDS Clean up
STEP 4: INJECT MEDICATION INTO THE DEVICE
Locate the cap. ❖ Open an alcohol pad and place the pad on your workspace. ❖ Locate the white cap on the top of the eclipse device. ❖ Remove the cap and place it on the alcohol pad.
Attach the syringe to the top of the eclipse device by screwing the syringe onto the eclipse. Slowly inject the syringe contents into the eclipse device.
Remove the syringe from the eclipse by unscrewing it and setting it aside. Replace the white cap on the eclipse device.
Discard all used supplies and trash in the garbage.
 WASH YOUR HANDS. Soap and water are preferred, hand sanitizer optional if soap and water are not available. Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.

Disclosure: This project was funded under grant number 1K08HS025782-01 from the Agency for Healthcare Research and Quality (AHRQ), U.S. Department of Health and Human Services (HHS). The authors are solely responsible for this document's contents, findings, and conclusions, which do not necessarily represent the views of AHRQ. Readers should not interpret any statement in this report as an official position of AHRQ or of HHS. None of the authors has any affiliation or financial involvement that conflicts with the material presented in this report.

