Practice good hand-washing

INSTRUCTIONAL CHECKLIST:

Titrate Up and Go – How To Discontinue Your IVIG Infusion

Prepare supplies

https://www.youtube.com/watch?time_continue=2&v=0im2OWCjM_g&feat ure=emb_logo

For additional education please contact your agency for assistance

	Four Ste	eps for	Perforr	ning I	nfusions:
--	----------	---------	---------	--------	-----------

2

Disconnect and flush

Medication infused:		Home Infusion Nurse (name, #):					
		Home Infusio	on Pharma	acist (name,	#):		
	ays wash hands Sc nen instructed	pap and water preferred		Hand sanitize (optional)		Wash for a conds or rub	
	STEP 1: PRACT	TICE GOOD	HAND-\	WASHING			
 Soap and wat Make it a hab If using hand hands to spee Remember, if 	to WASH YOUR HANDS for ter is always preferred. If so bit to wet your hands, apply sanitizer, rub a dime-sized a ed up drying. If you are using gloves, you re thands helps reduce your c	oap and water a soap, and lath amount betwee must still wash	rea not aver for 20 sen your ha	seconds (hur ands until dr	n the Happy	Birthday so	ong twice).
Prepare supplies Ass	sess device WASH HANDS	Gather supp	olies	WASH HANDS	Arrang supplie	—	WASH HANDS
	STEP 2	2: PREPARE	SUPPLII	ES			
program completing is alarming for anot	een titrated to the maximg or the infusion bag/both her reason, you should hing until the pump beeps a	tle running dr it the STOP bu	y, you sho	ould follow	the steps b	elow. If yo	our infusion
□ WASH YOUR HAND	S.		1			1.	



Discontinue the IV

	Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
	Clean your workspace using a cleaning solution (alcohol/sanitizing wipes or Lysol + paper towels).
	Select a dry, hard work surface to set up your supplies.
	WASH YOUR HANDS.
	 Soap and water are preferred (hand sanitizer optional). Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
	Collect supplies:
	 1 Alcohol Pad (to clean your end cap) 1 Saline Syringe (to clear your IV line and remove any leftover medication) 1 2X2 gauze and a band aid if desired 1 Trash bag
	Arrange your alcohol pad, wrapped syringes, and gauze above your SASH mat. You are at the last S and will not need the H to complete the administration. Do <u>not</u> place any of these items on top of your SASH mat yet. Keep the plastic bag close for trash disposal.
	 "S" = Saline "A" = Administer or Antibiotic (includes your medication bag, gravity tubing and red cap) "S" = Saline "H" = Heparin
	Prepare your supplies by unwrapping the saline syringe and setting aside. Check the unwrapped syringes for any air bubbles. If there are air bubbles follow the steps below.
	 Remove the protective cap from the syringe and discard. Hold the syringe with the end up and plunger at the bottom. Pull back on the syringe plunger slightly to break the seal. You will be able to see the air bubble in the syringe. Slowly push the plunger to let the air bubble out. Once the air is removed, set aside.
Disco	Disconnect and WASH HANDS Disconnect tubing WASH HANDS Scrub the Hub Pulsatile flush
	STEP 3: DISCONNECT AND FLUSH
	WASH YOUR HANDS.
	 Soap and water are preferred, hand sanitizer is optional if soap and water are not available. Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
	Stop the pump and disconnect.
	 When the pump alarms indicating the infusion has completed, Stop the pump by hitting the Red/Green Stop/Start button on the device. Select YES when asked "Stop Pump". Turn the pump off by pressing and holding down the Power Switch on the side of the pump. You will now disconnect your completed infusion from your IV line. Take the blue end cap (hub) of your IV line with one hand while using your other hand to unscrew the medication tubing. Do this until it is disconnected from the hub. Do not remove the end cap from the hub of your IV line.



	and be sure to hold the blue end cap of your IV and not the tubing. Unscrew the tubing careful to not remove the blue end cap from the hub of the vascular access device tubing.
	Remove the used cassette and discard. Make sure the clamp is closed on the tubing. If the cassette is locked, Insert the pump key into the key slot located on the side of the pump and turn counter clockwise into the unlocked position. Your cassette may not be locked and if not that is ok. Push down on the latch located on the side of the pump until the cassette detaches from the pump. Discard of the used cassette and bag in the trash bag.
	WASH YOUR HANDS.
	 Soap and water are preferred, hand sanitizer is optional if soap and water are not available. Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
	'Scrub the hub' of your IV line with an alcohol pad for at least 15 seconds.
	 Use an alcohol pad to scrub the end of the IV tubing. Once you have cleaned the hub, hold the hub with two fingers. Keep the hub from touching anything else. If your hub touches anything, wash your hands and scrub the hub again with a new alcohol pad.
	 Flush your IV line with saline ("pulse flush"): Hold the hub in one hand. Grasp the saline syringe on the first "S" of your SASH mat with your other hand. Remove the protective cover on the saline syringe. Screw the syringe onto the hub using a "push and twist" motion. It's like putting a key into an ignition. Once the syringe is connected to the hub, open the clamp on your IV line and slowly push 1 to 2 mL of saline at a time. Stop and push 1 to 2 mL more. Repeat this "pulse flush" until the saline syringe is empty. Flushing this way each time helps keep your IV line from getting clogged. If you have trouble flushing your IV line, make sure the clamp on your IV line is open. If your clamp is open and you are still having trouble, stop and call your home infusion nurse.
	Remove the empty saline syringe from your IV line. Throw it away.
	 WASH YOUR HANDS. ❖ Soap and water are preferred, hand sanitizer is optional if soap and water are not available. ❖ Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
Disco	ontinue the IV Remove the dressing Slowly remove Apply pressure Apply pressure Place bandaid WASH HANDS
	STEP 4: DISCONTINUE THE IV
	You will remove the peripheral vascular access device only if you have been directed to by your nurse. If you are unsure, do not remove the IV without contacting your IV company.
	Slowly peel back the tape from the IV line careful to not remove the IV with the tape/dressing. Discard in the trash.



Remove the catheter:
 Open the gauze and remove it from the packaging. Grasp the catheter by the hub and slowly but firmly pull the catheter away from the patient. As you remove the catheter use the gauze to apply pressure to the insertion site. Hold pressure for a minute and check for ongoing bleeding. Hold pressure until bleeding has stopped. Place Band-Aid if desired over the insertion site.
WASH YOUR HANDS.
 Soap and water are preferred, hand sanitizer is optional if soap and water are not available. Wash for 20 seconds (hum the Happy Birthday song twice) or rub with hand sanitizer until your hands are dry.
Congratulations! You have successfully disconnected and removed your vascular access device.
Make a habit to: □ WASH YOUR HANDS for at LEAST 20 seconds. ❖ Soap and water preferred (use hand sanitizer if soap or water are unavailable). ❖ Hum the Happy Birthday song twice. □ Contact your care team in the event you are not sure if the IV it to be removed. □ Always be sure that your medication has finished infusing by checking to see what is left in the container. Call your nurse if you have questions. □ Don't forget if you begin to feel itching, notice swelling, or other symptoms, call 911 and administer your EPI Pen as instructed.

Disclosure: This project was funded under grant number 1K08HS025782-01 from the Agency for Healthcare Research and Quality (AHRQ), U.S. Department of Health and Human Services (HHS). The authors are solely responsible for this document's contents, findings, and conclusions, which do not necessarily represent the views of AHRQ. Readers should not interpret any statement in this report as an official position of AHRQ or of HHS. None of the authors has any affiliation or financial involvement that conflicts with the material presented in this report.

