



JHCP VAXELIS SCHEDULE

	ROUTINE	SEASONAL
2 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM
	PREVNAR20 (PCV20)	0.5 ml IM
	ROTATEQ (rotavirus)	2 ml Oral
4 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM
	PREVNAR20 (PCV20)	0.5 ml IM
	ROTATEQ (rotavirus)	2 ml Oral
6 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM
	PREVNAR20 (PCV20)	0.5 ml IM
	ROTATEQ (rotavirus)	2 ml Oral
12 MO	MMR II (measles, mumps, rubella)	0.5 ml IM
	VARIVAX (varicella)	0.5 ml IM
	VAQTA (Hep A)	0.5 ml IM
15 MO	DAPTACEL (DTaP)	0.5 ml IM
	PEDVAXHIB (Hib)	0.5 ml IM
	PREVNAR20 (PCV20)	0.5 ml IM
18 MO	VAQTA (Hep A) *6 mo after first dose	0.5 ml IM

NB
-
8 mo
(1 time dose)
BEYFORTUS
(rsv)

< 5 kg
= 0.5 ml IM
≥ 5 kg
= 1 ml IM

6 mo+
(annually)
Influenza (flu)
vaccines
&
COVID vaccine

For flu and
covid, give the
age appropriate
formulation and
dose/s per
current CDC
guidance and
patient
vaccination
history

This guide reflects routine CDC recommended pediatric vaccines with the brands on the current JHCP formulary. Children with high risk conditions may receive addition vaccines or doses based on CDC recommendations.



JHCP VAXELIS SCHEDULDE

ROUTINE		SEASONAL
4Y - 6Y	QUADRACEL (DTaP, IPV-polio)	0.5 ml IM
	PROQUAD (MMR, varicella)	0.5 ml IM
9Y	GARDASIL 9 (human papilloma virus)	0.5 ml IM
	Give dose #2, 6 months-12 months after dose #1	
11Y	GARDASIL 9 (human papilloma virus)	0.5 ml IM
	BOOSTRIX (Tdap)	0.5 ml IM
16Y	MENQUADFI (meningococcal ACYW)	0.5 ml IM
	MENQUADFI (meningococcal ACYW)	0.5 ml IM
	BEXSERO (meningococcal b)	0.5 ml IM
	Give Bexsero dose #2, 1 month after dose #1	
	BEXSERO (meningococcal b)	0.5 ml IM
	For flu and covid, give the age appropriate formulation and dose/s per current CDC guidance and patient vaccination history	
6 mo+ (annually) Influenza (flu) vaccines & COVID vaccine		

This guide reflects routine CDC recommended pediatric vaccines with the brands on the current JHCP formulary. Children with high risk conditions may receive addition vaccines or doses based on CDC recommendations.