

VAXELIS SCHEDULE

ROUTINE			SEASONAL
2 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM	<p>NB-8 mo</p> <p>(1 time dose if mother did not received Abrysvo >2 weeks prior to delivery)</p> <p>BEYFORTUS (rsv)</p> <p>< 5 kg = 0.5 ml IM</p> <p>≥ 5 kg = 1 ml IM</p> <p>6 mo+ (annually)</p> <p>Influenza (flu) vaccines</p> <p>&</p> <p>COVID vaccine</p> <p>For flu and covid, give the age appropriate formulation and dose/s per current CDC guidance and patient vaccination history</p>
	PREVNAR20 (PCV20)	0.5 ml IM	
	ROTATEQ (rotavirus)	2 ml Oral	
4 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM	
	PREVNAR20 (PCV20)	0.5 ml IM	
	ROTATEQ (rotavirus)	2 ml Oral	
6 MO	VAXELIS (Hib, DTaP, IPV-Polio, Hep B)	0.5 ml IM	
	PREVNAR20 (PCV20)	0.5 ml IM	
	ROTATEQ (rotavirus)	2 ml Oral	
12 MO	MMR II (measles, mumps, rubella)	0.5 ml IM	
	VARIVAX (varicella)	0.5 ml IM	
	VAQTA (Hep A)	0.5 ml IM	
15 MO	DAPTACEL (DTaP)	0.5 ml IM	
	PEDVAXHIB (Hib)	0.5 ml IM	
	PREVNAR20 (PCV20)	0.5 ml IM	
18 MO	VAQTA (Hep A) *6 mo after first dose	0.5 ml IM	

This guide reflects routine CDC recommended pediatric vaccines with the brands on the current JHCP formulary. Children with high risk conditions may receive addition vaccines or doses based on CDC recommendations.

VAXELIS SCHEDULE

ROUTINE			SEASONAL
4Y - 6Y	QUADRACEL (DTaP, IPV-polio)	0.5 ml IM	6 mo+ (annually) Influenza (flu) vaccines & COVID vaccine For flu and covid, give the age appropriate formulation and dose/s per current CDC guidance and patient vaccination history
	PROQUAD (MMR, varicella)	0.5 ml IM	
9Y	GARDASIL 9 (human papilloma virus)	0.5 ml IM	
	Give dose #2, 6 months-12 months after dose #1		
	GARDASIL 9 (human papilloma virus)	0.5 ml IM	
11Y	BOOSTRIX (Tdap)	0.5 ml IM	
	MENVEO (meningococcal ACYW)	0.5 ml IM	
16Y	MENVEO (meningococcal ACYW)	0.5 ml IM	
	BEXSERO (meningococcal b)	0.5 ml IM	
	Give Bexsero dose #2, 6 months after dose #1		
	BEXSERO (meningococcal b)	0.5 ml IM	

This guide reflects routine CDC recommended pediatric vaccines with the brands on the current JHCP formulary. Children with high risk conditions may receive addition vaccines or doses based on CDC recommendations.