

If you have any questions, regarding the effect of any medications on the list, please call our office @ 410-760-8840.

Please do not take any of the following **48 hours** before the test. However, **contact your physician should you have any reservations about discontinuing any medications.**

Antibiotics:	Gentamicin, Streptomycin
Anticonvulsants:	Klonopin (do not stop taking for anxiety), Dilantin, Librium, Librax, Tegretol (carbamazepine), Nembutal
Antidepressants/Sedatives:	Ativan, Elavil, Lithium, Pamelor, Placidyl, Prozac, Valium, Xanax
Antihistamines/Decongestants:	Allegra, Benadryl, Chlor-Trimeton, Claritin, Seldane, Sudafed, Zyrtec, Xyzal, Over-the-Counter cold remedies
Anti-nausea medicine:	Dramamine, Compazine, Bonine, Marezine, Votrol, Phenergan, Thorazine, etc.
Anti-vertigo medicine:	Anti-vert, Meclizine, Ru-vert, Transderm Scop patches, Vontrol Transderm Scop patches - must be off for 5 days prior to testing
Narcotics and Barbituates:	Phenobarbital, Codeine, Demerol, Dilaudid, Percocan, Percocet, Phenaphen, etc.
Sleeping Medications:	Ambien, Dalmane, Doral, Halcion, Lunesta, ProSom, Restoril, Seconal, Unisom
Stimulant Medications:	Amphetamine, Ritalin
Tranquilizers:	Valium, Librium, Atarax, Visteril, Equanil, Miltown, Travil, Serex, Etrafon, Ativan, etc.
Other Medications:	Alcohol (beer, wine, liquor, or cough medications containing alcohol) Methadone, NSAID, Quinine, Salicylates-Aspirin

No recreational drugs

** Anti-Epilepsy and Anti-Psychotic medications, as well as medications for Diabetes, Hypertension, or Heart Disease **should not be discontinued.**

Annapolis

Columbia

Glen Burnie

Kent Island

Laurel

Odenton