

STANDARD MEDICARE PART B MANAGEMENT

**REMICADE (infliximab)
AVSOLA (infliximab-axxq)
INFLECTRA (infliximab-dyyb)
RENFLEXIS (infliximab-abda)
infliximab**

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Crohn's disease
 - i. Reducing signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active Crohn's disease (CD) who have had an inadequate response to conventional therapy.
 - ii. Reducing the number of draining enterocutaneous and rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing CD.
2. Pediatric Crohn's disease
Reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients 6 years of age and older with moderately to severely active CD who have had an inadequate response to conventional therapy.
3. Ulcerative colitis
Reducing signs and symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active ulcerative colitis (UC) who have had an inadequate response to conventional therapy.
4. Pediatric ulcerative colitis
Reducing signs and symptoms and inducing and maintaining clinical remission in pediatric patients 6 years of age and older with moderately to severely active UC who have had an inadequate response to conventional therapy.
5. Rheumatoid arthritis in combination with methotrexate
Reducing signs and symptoms, inhibiting the progression of structural damage, and improving physical function in patients with moderately to severely active rheumatoid arthritis (RA).
6. Ankylosing spondylitis
Reducing signs and symptoms in adult patients with active ankylosing spondylitis (AS).
7. Psoriatic arthritis
Reducing signs and symptoms of active arthritis, inhibiting the progression of structural damage, and improving physical function in adult patients with psoriatic arthritis (PsA).
8. Plaque Psoriasis

Reference number
1979-A

Treatment of adult patients with chronic severe (i.e., extensive and/or disabling) plaque psoriasis (PsO) who are candidates for systemic therapy and when other systemic therapies are medically less appropriate.

B. Compendial Uses

1. Adult-onset Still's disease
2. Arthritis in Crohn's disease
3. Axial spondyloarthritis
4. Behcet's disease
5. Gastrointestinal tract transplantation organ rejection
6. Giant cell arteritis
7. Acute graft versus host disease
8. Hidradenitis suppurativa
9. Juvenile idiopathic arthritis
10. Kawasaki disease
11. Necrobiosis lipoidica diabetorum
12. Pyoderma gangrenosum
13. Rheumatoid arthritis as monotherapy
14. Severe, refractory SAPHO (synovitis, acne, pustulosis, hyperostosis, and osteitis) syndrome
15. Sarcoidosis
16. Subcorneal pustular dermatosis
17. Synovitis
18. Takayasu's arteritis
19. Uveitis
20. Immune checkpoint inhibitor toxicity
21. Multisystem inflammatory syndrome in children (MIS-C)

All other indications will be assessed on an individual basis. Submissions for indications other than those enumerated in this policy should be accompanied by supporting evidence from Medicare approved compendia.

II. DOCUMENTATION

The following documentation must be available, upon request, for all submissions:

- A. Crohn's disease (CD) and ulcerative colitis (UC)
For continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.
- B. Rheumatoid arthritis (RA)
 1. For initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 2. For continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- C. Ankylosing spondylitis (AS), axial spondyloarthritis (axSpA), psoriatic arthritis (PsA), adult-onset Still's disease (AOSD), hidradenitis suppurativa, juvenile idiopathic arthritis (JIA), and uveitis

Reference number
1979-A

For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

- D. **Plaque psoriasis (PsO)**
For continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.
- E. **Acute graft versus host disease (initial requests only)**
Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

III. CRITERIA FOR INITIAL APPROVAL

- A. **Crohn's disease (CD)**
Authorization of 12 months may be granted for treatment of moderately to severely active Crohn's disease.
- B. **Ulcerative colitis (UC)**
Authorization of 12 months may be granted for treatment of moderately to severely active ulcerative colitis.
- C. **Rheumatoid arthritis (RA)**
Authorization of 12 months may be granted for treatment of moderately to severely active rheumatoid arthritis when any of the following criteria is met:
 1. The member has previously received any other biologic drug or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for rheumatoid arthritis.
 2. The member had an inadequate response to methotrexate or leflunomide.
 3. The member has a clinical reason to avoid treatment with methotrexate or leflunomide (e.g., renal or hepatic impairment, pregnancy or currently planning pregnancy).
- D. **Ankylosing spondylitis (AS) and axial spondyloarthritis (axSpA)**
Authorization of 12 months may be granted for treatment of active ankylosing spondylitis and active axial spondyloarthritis.
- E. **Psoriatic arthritis (PsA)**
Authorization of 12 months may be granted for treatment of active psoriatic arthritis.
- F. **Plaque psoriasis (PsO)**
Authorization of 12 months may be granted for treatment of plaque psoriasis.
- G. **Adult-onset Still's disease (AOSD)**
Authorization of 12 months may be granted for treatment of active adult-onset Still's disease.
- H. **Arthritis in Crohn's disease (CD)**
Authorization of 12 months may be granted for treatment of arthritis in a member with Crohn's disease.
- I. **Behcet's disease**
Authorization of 12 months may be granted for treatment of Behcet's disease.

Reference number
1979-A

J. Gastrointestinal tract transplantation organ rejection

Authorization of 6 months may be granted for treatment of gastrointestinal tract transplantation organ rejection.

K. Giant cell arteritis

Authorization of 3 months may be granted for treatment of giant cell arteritis.

L. Acute graft versus host disease

Authorization of 12 months may be granted for treatment of acute graft versus host disease when either of the following criteria is met:

1. Member has experienced an inadequate response to systemic corticosteroids.
2. Member has an intolerance or contraindication to corticosteroids.

M. Hidradenitis suppurativa

Authorization of 12 months may be granted for treatment of hidradenitis suppurativa.

N. Juvenile idiopathic arthritis (JIA)

Authorization of 12 months may be granted for treatment of active juvenile idiopathic arthritis.

O. Kawasaki disease

Authorization of 1 month may be granted for treatment of Kawasaki disease.

P. Necrobiosis lipoidica diabetorum

Authorization of 12 months may be granted for treatment of necrobiosis lipoidica diabetorum.

Q. Pyoderma gangrenosum

Authorization of 12 months may be granted for treatment of pyoderma gangrenosum.

R. SAPHO (synovitis, acne, pustulosis, hyperostosis, and osteitis) syndrome

Authorization of 12 months may be granted for treatment of severe, refractory SAPHO syndrome.

S. Sarcoidosis

Authorization of 12 months may be granted for treatment of sarcoidosis.

T. Subcorneal pustular dermatosis

Authorization of 6 months may be granted for treatment of subcorneal pustular dermatosis.

U. Synovitis

Authorization of 12 months may be granted for treatment of synovitis.

V. Takayasu's disease

Authorization of 12 months may be granted for treatment of Takayasu's disease.

W. Uveitis

Authorization of 12 months may be granted for treatment of uveitis.

X. Immune checkpoint inhibitor toxicity

Authorization of 6 months may be granted for treatment of immune checkpoint inhibitor toxicity when the member meets either of the following criteria:

Reference number
1979-A

1. Member has not responded to systemic corticosteroids.
2. Member has moderate or severe diarrhea or colitis.

Y. Multisystem inflammatory syndrome in children (MIS-C)

Authorization of 1 month may be granted for treatment of multisystem inflammatory syndrome in children (MIS-C) post severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection who have failed to respond to standard pharmacologic therapy.

IV. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must be currently receiving therapy with the requested agent.

A. Crohn's disease (CD)

Authorization for 12 months may be granted for moderately to severely active Crohn's disease when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as one of the following:
 - i. Member has achieved or maintained remission.
 - ii. Member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - a. Abdominal pain or tenderness
 - b. Diarrhea
 - c. Body weight
 - d. Abdominal mass
 - e. Hematocrit
 - f. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
 - g. Improvement on a disease activity scoring tool (e.g., Crohn's Disease Activity Index [CDAI] score)

B. Ulcerative colitis (UC)

Authorization for 12 months may be granted for moderately to severely active ulcerative colitis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as one of the following:
 - i. Member has achieved or maintained remission.
 - ii. Member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - a. Stool frequency
 - b. Rectal bleeding
 - c. Urgency of defecation
 - d. C-reactive protein (CRP)
 - e. Fecal calprotectin (FC)
 - f. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound

Reference number
1979-A

- g. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

C. Rheumatoid arthritis (RA)

Authorization for 12 months may be granted for moderately to severely active rheumatoid arthritis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by disease activity improvement from baseline in tender joint count, swollen joint count, pain, or disability.

D. Ankylosing spondylitis (AS) and axial spondyloarthritis (axSpA)

Authorization for 12 months may be granted for active ankylosing spondylitis or active axial spondyloarthritis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Functional status
 - ii. Total spinal pain
 - iii. Inflammation (e.g., morning stiffness)

E. Psoriatic arthritis (PsA)

Authorization for 12 months may be granted for psoriatic arthritis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Number of swollen joints
 - ii. Number of tender joints
 - iii. Dactylitis
 - iv. Enthesitis
 - v. Axial disease
 - vi. Skin and/or nail involvement

F. Plaque psoriasis (PsO)

Authorization for 12 months may be granted for plaque psoriasis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. Member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when either of the following is met:
 - i. Reduction in body surface area (BSA) affected from baseline
 - ii. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

G. Adult-onset Still's disease (AOSD)

Authorization for 12 months may be granted for adult-onset Still's disease when both of the following criteria are met:

Reference number
1979-A

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
 - ii. Number of joints with limitation of movement
 - iii. Functional ability
 - iv. Systemic features (e.g., fever, evanescent rash, lymphadenopathy, hepatomegaly, splenomegaly, serositis)

H. Hidradenitis suppurativa

Authorization for 12 months may be granted for hidradenitis suppurativa when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:
 - i. Reduction in abscess and inflammatory nodule count from baseline
 - ii. Reduced formation of new sinus tracts and scarring
 - iii. Decrease in frequency of inflammatory lesions from baseline
 - iv. Reduction in pain from baseline
 - v. Reduction in suppuration from baseline
 - vi. Improvement in frequency of relapses from baseline
 - vii. Improvement in quality of life from baseline
 - viii. Improvement on a disease severity assessment tool from baseline

I. Active juvenile idiopathic arthritis (JIA)

Authorization for 12 months may be granted for active juvenile idiopathic arthritis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
 - ii. Number of joints with limitation of movement
 - iii. Functional ability

J. Uveitis

Authorization for 12 months may be granted for uveitis when both of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The member is receiving benefit from therapy. Benefit is defined as the member has achieved or maintained a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:
 - i. Reduced frequency of recurrence compared to baseline
 - ii. Zero anterior chamber inflammation or reduction in anterior chamber inflammation compared to baseline
 - iii. Decreased reliance on topical corticosteroids

Reference number
1979-A

K. Gastrointestinal tract transplantation organ rejection, giant cell arteritis, Kawasaki disease, subcorneal pustular dermatosis, and immune checkpoint inhibitor toxicity

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

L. All other indications

Authorization for 12 months may be granted when all of the following criteria are met:

1. The member is currently receiving therapy with Avsola, Inflectra, infliximab, Remicade, or Renflexis.
2. The requested medication is being used to treat an indication enumerated in Section III.
3. The member is receiving benefit from therapy.

V. SUMMARY OF EVIDENCE

The contents of this policy were created after examining the following resources:

1. The prescribing information for Remicade and its biosimilars.
2. The available compendium
 - a. National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium
 - b. Micromedex DrugDex
 - c. American Hospital Formulary Service- Drug Information (AHFS-DI)
 - d. Lexi-Drugs
3. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update
4. 2015 American College of Rheumatology guidelines for the treatment of rheumatoid arthritis
5. 2010 Update of the international ASAS recommendations for the use of anti-TNF agents in patients with axial spondyloarthritis
6. EULAR recommendations on management of Behcet's syndrome.
7. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations: Part II: Topical, intralesional, and systemic medical management.
8. British Association of Dermatologists guidelines for the management of hidradenitis suppurativa (acne inversa) 2018
9. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis.
10. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis.
11. Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals from the American Heart Association.
12. Etiology and management of pyoderma gangrenosum: a comprehensive review
13. European Respiratory Society (ERS) clinical practice guidelines on treatment of sarcoidosis.
14. Comparison of ultrasonographic assessment of synovitis and joint vascularity with radiographic evaluation in a randomized, placebo-controlled study of infliximab therapy in early rheumatoid arthritis.
15. Very early treatment with infliximab in addition to methotrexate in early, poor-prognosis rheumatoid arthritis reduces magnetic resonance imaging evidence of synovitis and damage, with sustained benefit after infliximab withdrawal: results from a twelve-month randomized, double-blind, placebo-controlled trial.

Reference number
1979-A

16. Recommendations of the Italian Society of Rheumatology for the treatment of the primary large-vessel vasculitis with biological agents
17. Efficacy and tolerance of infliximab in refractory Takayasu arteritis: French multicentre study.
18. A review of systemic biologics and local immunosuppressive medications in uveitis
19. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders.
20. NCCN guideline: Hematopoietic cell transplantation
21. NCCN guideline: Management of immunotherapy-related toxicities
22. COVID-19 Treatment Guidelines Panel: Coronavirus disease 2019 (COVID-19) treatment guidelines.

After reviewing the information in the above resources, the FDA-approved indications listed in the prescribing information for Remicade and its biosimilars are covered in addition to the following:

- A. Axial spondyloarthritis
- B. Adult-onset Still's disease
- C. Arthritis in Crohn's disease
- D. Behçet's disease
- E. Gastrointestinal tract transplantation organ rejection
- F. Giant cell arteritis
- G. Acute graft versus host disease
- H. Hidradenitis suppurativa
- I. Juvenile idiopathic arthritis
- J. Kawasaki disease
- K. Necrobiosis lipoidica diabetorum
- L. Pyoderma gangrenosum
- M. Sarcoidosis
- N. Subcorneal pustular dermatosis
- O. Synovitis
- P. Takayasu's arteritis
- Q. Uveitis
- R. Immune checkpoint inhibitor toxicity
- S. Multisystem inflammatory syndrome in children (MIS-C)

VI. EXPLANATION OF RATIONALE

Support for FDA-approved indications (Crohn's disease, ulcerative colitis, rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, and plaque psoriasis) can be found in the manufacturer's prescribing information.

For established RA, the guidelines state "if the disease activity is low, in patients who have never taken a DMARD, the recommendation is to use DMARD monotherapy (methotrexate preferred) over TNFi". "If disease activity remains moderate or high despite DMARD monotherapy, the recommendation is to use combination traditional [conventional] DMARDs or add a TNFi or a non-TNF biologic or tofacitinib (all choices with or without methotrexate, in no particular order of preference), rather than continuing DMARD monotherapy alone". Recommendations for patients with symptomatic early RA state that "if disease activity is low, in patients who have never taken a DMARD, use DMARD monotherapy (methotrexate preferred) over double or triple therapy". "If disease activity remains moderate or high despite DMARD monotherapy (with or without glucocorticoids), use combination DMARDs or a TNFi or a non-TNF biologic (all choices with or without methotrexate, in no particular order of preference), rather than continuing DMARD monotherapy alone". A strong recommendation means that the panel was confident that the desirable effects of following the

Reference number
1979-A

recommendation outweigh the undesirable effects (or vice versa), so the course of action would apply to most patients, and only a small proportion would not want to follow the recommendation (Singh et al., 2016).

Support for using infliximab for axial spondyloarthritis can be found in the 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. The guidelines recommend that patients who still have active AS despite treatment with NSAIDs, tumor necrosis factor inhibitor (TNFi) such as infliximab are recommended but the guideline does not recommend any particular TNFi.

Support for using infliximab to treat adult-onset Still's disease can be found in two published case series. Kraetsch et al reported adult-onset Still disease (AOSD) appears to favorably respond to treatment with infliximab. In a small, pilot study, 6 patients diagnosed with AOSD (4 with early onset of disease, and 2 with disease durations of 3 and 5 years, respectively) received an initial course of intravenous infusions of infliximab 5 mg/kg, at 0, 2, and 6 weeks. Further treatment with infliximab was given at 6- to 8-week intervals, contingent upon patient response. At the time of study enrollment, all patients had massive polyarthralgia, 5 had polyarthritis, 5 had persistent fever, 5 had a characteristic rash, 5 had persistent leukocytosis, 4 had splenomegaly, and all 6 patients had elevations of erythrocyte sedimentation rate (ESR) and elevated serum concentrations of C-reactive protein. Hyperferritinemia was seen in 3 patients. All patients showed a beneficial response to treatment, with complete resolution of rash, fever, myalgias, and splenomegaly (the latter after 3 treatments); arthralgia/arthritis resolved in 5 of 6 patients. Normalization also occurred in serological markers of disease activity (CRP, ESR, and ferritin concentration) in all patients. Favorable effects of treatment were evident after the first course of treatment with infliximab and were sustained with continuing infliximab treatment at 6-to-8 week intervals, with treatment durations extending from 5 to 28 months. In the 2 patients with long-standing disease of 3- and 5-years duration, swollen joint counts declined from 30 to 3, and from 3 to 0 joints, respectively; tender joint count declined similarly, from 33 to 3 and from 7 to 2 joints, respectively. Infliximab was tolerated well; 1 patient showed a moderate infusion reaction during the second treatment yet was able to resume infliximab therapy after a brief discontinuation of the infusion. Cavagna et al indicated that infliximab appeared to induce clinical remission in 3 patients with chronically active, treatment-refractory, adult Still Disease (ASD). Each patient had a disease history of between 4- and 7-years duration, during which time they exhibited relapsing or refractory disease despite treatment with NSAIDs, prednisone, methotrexate (n=3), and cyclosporine (n=1). Patients were given intravenous infusions of infliximab 3 mg/kg at weeks 0, 2, 6, and then once every 8 weeks. Infliximab was to be given once every 4 weeks from week 30 thereafter, and methotrexate was maintained throughout the duration of the study. All patients experienced rapid regression of ASD symptoms (arthralgia, cutaneous rash, fever, pharyngitis), accompanied by progressive reductions in serum concentrations of ferritin, C-reactive protein, and erythrocyte sedimentation rate. One patient developed a diffuse, urticarial rash shortly after the fifth infliximab infusion, necessitating withdrawal from therapy at week 22. The 2 remaining patients both experienced brief relapses on weeks 20 and 28; both rapidly regained a state of remission following repeat infusions of infliximab, and continued to receive infliximab beyond 30 weeks, without signs of relapse. These 2 patients also tolerated tapered reductions in prednisone dosing. Neither of the remaining patients showed development of anticardiolipin antibodies, anti-double stranded DNA, or antinuclear antibodies after prolonged treatment.

Support for using infliximab to treat arthritis in Crohn's disease can be found in a case series by Elman et al. Infliximab appeared to be effective in suppressing joint inflammation associated with arthritis secondary to Crohn disease. In a series of case reports, patients with treatment-refractory, joint inflammation associated with long-standing Crohn disease (9 to 31 years duration; n=4) were given intravenous infliximab 5 mg/kg at 8 to 16-week intervals after the initial induction schedule. All patients had prolonged episodes of joint and back pain associated with periods of quiescence in their inflammatory bowel disease; 2 patients had "sausage-

Reference number
1979-A

shaped" finger and toe swelling, and 3 patients presented with pain in the sacroiliac joint (SI-joint). No patient had radiographic abnormalities of the SI-joint. All patients had been receiving treatment with prednisone up to 40 mg per day, accompanied by 1 or more antiarthritic agents including azathioprine, methotrexate, minocycline, and sulfasalazine. Patients had favorable responses to infliximab, experiencing clinically meaningful reductions in joint pain and swelling, allowing for dose reductions or withdrawals of corticosteroid and antiarthritic agents. One patient discontinued infliximab therapy due to anorexia and insomnia.

Support for Behcet's disease can be found the European League Against Rheumatism (EULAR) recommendations on management of Behcet syndrome (BS). Hatemi and colleagues (2018) noted that several new therapeutic modalities with different mechanisms of action have been studied in patients with BS. These researchers updated the recommendations in the light of these new data under the auspices of EULAR Standing Committee for Clinical Affairs. The recommendations on the medical management of mucocutaneous, joint, eye, vascular, neurological and GI involvement of BS were modified; 5 overarching principles and a new recommendation about the surgical management of vascular involvement were added. For BS with eye involvement, among the monoclonal anti-TNF antibodies, although there is more accumulated experience with IFX, ADA also appeared to be an effective alternative. Switching between these agents appeared to be possible in patients with primary or secondary unresponsiveness or AEs. Patients presenting with an initial or recurrent episode of acute sight-threatening uveitis should be treated with high-dose glucocorticoids, IFX or IFN-alpha. Intravitreal glucocorticoid injection is an option in patients with unilateral exacerbation as an adjunct to systemic treatment.

Support for using infliximab to treat gastrointestinal tract organ transplantation rejection can be found in two case reports by Pascher et al. In 2 case reports, infliximab was effective in the treatment of steroid and OKT3 (muromonab-CD3)-refractory moderate to severe acute cellular rejection in intestinal transplant recipients. Following either 5 or 10 days of treatment with OKT3 and enhanced baseline immunosuppressive therapy, acute cellular rejection persisted. Patients were then treated with 3 mg/kg IV infliximab; both patients received 4 infusions, 2 to 4 weeks apart. Improvement was observed within 1 week of the first infusion. Absence of clinical symptoms and histological signs of rejection persisted for at least 8 months for 1 patient and at least 10 months for the other.

Support for using infliximab to treat giant cell arteritis can be found in an open-label case study (Cantini et al). Administration of infliximab was effective in provoking remission in patients with active, steroid-dependent giant cell arteritis (GCA). In an open-label case study, 4 patients with long-standing GCA (disease duration ranging from 42 to 54 months) were unable to tolerate the tapering of their daily corticosteroid dose to less than 12.5 mg. They were given a 3-dose regimen of intravenous infliximab 3 mg/kg, at 0, 2, and 6 weeks, concurrent with reduction of their steroid dose to prednisone 5 mg per day. Three patients experienced a complete response to infliximab therapy, exhibiting both clinical and humeral evidence of remission (resolution of cranial and systemic symptoms, articular symptoms, visual symptoms, and normalization of erythrocyte sedimentation rate and serum concentration of C-reactive protein) after the second dose of infliximab. These responders remained in remission for up to 6 months after the third infliximab infusion, without requiring further treatment with corticosteroid. The fourth patient initially showed a partial response to the first infusion; however, she experienced clinical relapse at the time of her second infusion, causing her to withdraw from the study per the prospectively established protocol. Infliximab was well tolerated by all patients, and adverse events were neither reported nor observed.

Support for hidradenitis suppurativa can be found in the North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations: Part II: Topical, intralesional, and systemic medical management. The guideline indicates

Reference number
1979-A

infliximab is recommended for moderate-to-severe disease. Dose ranging studies are needed to determine the optimal dosage for management.

Support for juvenile idiopathic arthritis can be found in the 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Entesitis. Using infliximab in combination with a DMARD was a strong recommendation despite the low quality of evidence, primarily given more extensive experience with the need for combination therapy to reduce the risk of antidrug antibody formation.

Support for Kawasaki disease can be found in the following document produced by the American Heart Association: Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals from the American Heart Association. A 2-center, randomized, double-blind, placebo-controlled trial of infliximab plus IVIG for intensification of initial treatment enrolled 196 subjects. The study was powered for the primary outcome measure of reducing IVIG resistance from 20% to 5%. Secondary outcome measures included reduction of inflammatory parameters and the change in coronary artery Z scores. Although the number of fever days was shortened and inflammatory parameters normalized more rapidly in the infliximab-treated subjects, the rates of IVIG resistance were identical between the 2 arms. A striking finding was the complete prevention of IVIG infusion reactions in children randomized to the infliximab arm compared with a 13% reaction rate in subjects who received placebo before their IVIG infusion. There was a significant decrease in Z score for the LAD in favor of infliximab. However, there was no difference in the rate of coronary artery aneurysms between the groups, although the study was inadequately powered for this end point. On the basis of current information, addition of infliximab to initial therapy with IVIG is safe but does not prevent recrudescence fever.

A phase I multicenter, randomized, open clinical trial of infliximab (5 mg/kg intravenously over 2 hours) versus a second infusion of IVIG (2 g/kg) was performed to determine the safety, tolerability, and pharmacokinetics of infliximab for rescue therapy for patients who had fever at least 36 hours after the end of the initial IVIG infusion. The study enrolled 24 subjects with IVIG-resistant KD and determined that infliximab was well tolerated in infants and children with KD and that the pharmacokinetics were similar to adults, with circulating levels of the monoclonal antibody detected out to 10 weeks. In the Japanese trial, 20 KD patients resistant to 2 consecutive IVIG infusions (2 g/kg each) were treated with infliximab (5 mg/kg), and an apparent clinical response was achieved in 18 (90%). The 2 unresponsive patients were treated with plasma exchange with resolution of their inflammation. The coronary artery abnormalities detected by echocardiogram all subsequently resolved. There were no adverse reactions attributed to infliximab among the study subjects.

A retrospective review of 2 centers that consistently administered either a second dose of IVIG or infliximab to IVIG-resistant patients suggested that patients receiving infliximab had shorter hospitalization and fewer days of fever, but coronary artery outcomes and adverse events were similar. On the basis of these retrospective data, infliximab can be considered as an alternative to a second infusion of IVIG for resistant patients.

Support for using infliximab to treat necrobiosis lipoidica diabetorum can be found in a case report by Kolde et al. Infliximab was an effective treatment for refractory ulcerated necrobiosis lipoidica in a 33-year-old man with diabetes mellitus. The patient received once monthly infusions of infliximab (5 mg/kg) for 2 months. Following treatment with infliximab, clinical improvement was reported, including healing of ulcerations, fading of erythematous infiltration, flattening of the raised margin, and substantial reduction in pain. Improvement of the necrobiosis lipoidica was sustained after the cessation of infliximab. The only reported adverse event was the development of miliary tuberculosis after the second infusion, which was possibly drug-related due to the temporal association with infliximab treatment.

Reference number
1979-A

Support for pyoderma gangrenosum (PG) can be found in a study by Ahronowitz et al. Infliximab, an anti-TNF α monoclonal antibody binding both soluble and membrane-bound TNF α , is the only biologic that has shown efficacy in classic PG in a randomized, double-blind, controlled trial (level I evidence). Thirty patients were given either infliximab 5 mg/kg or placebo. At 2 weeks, 6 of 13 patients in the infliximab group showed improvement in the severity and/or size of ulcers, versus only 1 of 17 in the placebo group. After 2 weeks, the 16 non-responders in the placebo group were switched to infliximab and by week 6, 20 of 29 patients treated with infliximab demonstrated improvement in their PG lesions, with 6 of 29 showing complete resolution. Further studies are needed to determine the efficacy of infliximab in idiopathic PG.

Support for sarcoidosis can be found in the practice guidelines from the European Respiratory Society. The practice guidelines recommend the addition of infliximab to improve and/or preserve forced vital capacity (FVC) and quality of life in patients with symptomatic pulmonary sarcoidosis believed to be at higher risk for future mortality or permanent disability from sarcoidosis who have been treated with glucocorticoids or other immunosuppressive agents and have continued disease. Additionally, the guideline recommends the addition of infliximab (compared to no additional treatment) for patients with cutaneous sarcoidosis who have been treated with glucocorticoids and/or other immunosuppressive agents and have continued cosmetically important active skin disease. In patients with neurosarcoidosis that have been treated with glucocorticoids and a second-line agent (methotrexate, azathioprine, mycophenolate mofetil) with continued disease, the guidelines suggest adding infliximab.

Support for using infliximab to treat subcorneal pustular dermatosis can be found in a case report by Voightlander et al. Infliximab was effective in producing remission in a 79-year-old woman with treatment-refractory subcorneal pustular dermatosis (Sneddon-Wilkinson disease). The patient presented with disease flare (progressive, widespread erythema and pustular eruptions on the legs, forearms, trunk, and abdomen) that was recalcitrant to treatment with acitretin and methylprednisolone. Intravenous infliximab 5 mg/kg was given as a 2-hour infusion. Within 24 hours of treatment, serum analysis revealed a rapid decline in the number of peripheral granulocytes, accompanied by a decline (to within normal limits) in concentration of C-reactive protein. Complete resolution of pustules occurred within 2 days of infusion, leaving a residual scaling of the affected skin. The patient was able to tolerate the withdrawal of methylprednisolone over 3 days. Disease flare occurred 12 days after the first dose of infliximab; a second infliximab infusion (5 mg/kg) was given, provoking a complete remission within a day of the second treatment. Other than a mild, corticosteroid-responsive relapse, the patient remained in complete remission for a minimum of 6 months while receiving a maintenance therapy regimen of acitretin.

Support for synovitis can be found in a randomized, double-blind, placebo-controlled trial (n=20), significant reductions from baseline in MRI-measured synovitis were seen at 14 weeks and 1 year with infliximab plus methotrexate therapy. Disease Modifying Antirheumatic Drug- or oral corticosteroid-1 rheumatoid arthritis patients with recent symptom onset (less than 12 months) and with metacarpophalangeal joint involvement were randomized to methotrexate 7.5 mg once weekly plus infliximab 3 mg/kg or placebo infusion at 0, 2, 6, and then every 8 weeks. MRI-measured synovitis at week-14 (primary endpoint) and at week-54 (secondary endpoint) from baseline were compared between the infliximab and placebo groups. At week 14, median total synovitis score was significantly lower in the infliximab group (5.5 to 3.4) as compared with the placebo arm (6.2 to 5.9) (p less than 0.05). After 54 weeks, median total synovitis score was significantly lower in the infliximab group (3.8) as compared with the placebo arm (6.6) (p less than 0.05). Adverse effects with infliximab included infusion reaction (n=1), elevated liver function enzyme (n=1), and cutaneous vasculitis (n=1).

Additionally, in a placebo-controlled trial (n=24), infliximab plus methotrexate showed a significant percent reduction in total synovial thickness from baseline at 18 weeks compared with methotrexate alone. Patients with early phase rheumatoid arthritis (less than 2 years mean duration) with a minimum of 2 swollen

Reference number
1979-A

metacarpophalangeal joints despite methotrexate treatment were randomized to methotrexate (escalating dose) plus infliximab 5 mg/kg or placebo infusion at 0, 2, 6, and then every 8 weeks. After 18 weeks, high frequency ultrasonography showed a 50% median reduction in synovial thickness with the infliximab group as opposed to a 1.2% median increase in synovial thickness in the placebo group (p=0.014).

Support for Takayasu's arteritis can be found in the Recommendations of the Italian Society of Rheumatology for the treatment of the primary large-vessel vasculitis with biological agents. According to the recommendations, tumor necrosis factor alpha-inhibiting agents are recommended in patients with persistently active Takayasu Arteritis for 6 months or more, or with 2 or more flares or relapses despite glucocorticoid therapy; this is in addition to 1 or more immunosuppressive agent unless not tolerated or contraindicated. In a 12-month, multicenter, retrospective study (n=15), infliximab therapy resulted in a response rate of 73% to 87% and significantly reduced corticosteroid use in patients with refractory Takayasu arteritis. Patients (median age, 41 years; range, 17 to 61 years) with Takayasu arteritis (median time from disease onset to infliximab therapy, 37 months; range, 6 to 365 months) that was refractory to other nonsteroid immunosuppressive agents or steroids received infliximab 3 mg/kg (n=5) or 5 mg/kg (n=10) IV every 4 to 8 weeks (median, every 6 weeks). Patients were concomitantly receiving steroids (n=14; median prednisone dose, 20 mg; range, 5 to 35 mg/day) and other nonsteroid immunosuppressive therapies (methotrexate, n=7; azathioprine, n=4) with doses that were not modified in the 3 months before infliximab initiation. After a median follow-up of 43 months (range, 4 to 71 months), overall response (including partial or good response; determined by physician in change and by the presence of clinical and biological activity) was achieved in 87% (n=13/15), 77% (n=10/13), and 73% (n=8/11), respectively, at 3, 6, and 12 months. The percentage of patients with disease activity was significantly decreased from 73% at baseline to 20% at 3 months (p less than 0.005), 31% at 6 months (p less than 0.05), and 27% at 12 months (p less than 0.05). The median prednisone dose also significantly decreased from 20 mg (range, 5 to 35 mg) at baseline to 15 mg (range, 5 to 20 mg) at 3 months (p less than 0.005), 7.5 mg (range, 5 to 18 mg) at 6 months (p less than 0.05), and 6 mg (range, 2.5 to 30 mg) at 12 months (p less than 0.05). Additionally, C-reactive protein was decreased from a median of 30 mg/L (range, 4 to 70 mg/L) at baseline to 5 mg/L (range, 0 to 57 mg/L) at 3 months (p less than 0.05) and 6 mg/L (range, 0 to 50 mg/L) at 6 months (p less than 0.05); however, there was no significant difference from baseline at month 12. Adverse events included acute infusion reactions in 2 patients that led to discontinuation of infliximab therapy.

In a single center retrospective study (n=25), partial or complete remission occurred in 18 of 21 patients who received infliximab therapy for the treatment of refractory Takayasu arteritis. Patients (mean age, 35 years; range, 15 to 64 years; median disease duration, 116 months; range, 39 to 344 months; concurrent nonsteroid immunosuppressive therapy, n=18) with Takayasu arteritis who could not achieve stable remission with the use of low-dose prednisone (less than 10 mg/day) and who had received at least 1 additional immunosuppressive agent received infliximab (n=21) or etanercept (n=9). Five patients who were initially treated with etanercept were switched to infliximab. After a median follow-up of 28 months (range, 2 to 84 months), infliximab therapy (median dose 5 mg/kg IV (range, 4 to 10 mg/kg) every 6 weeks (range, 4 to 8 weeks)) resulted in remission (primary endpoint) in 18 patients (complete remission, n=12; partial remission, n=6). In patients who received either etanercept or infliximab, the median prednisone dose was 19 mg (range, 5 to 50 mg) prior to therapy compared with 0 mg (range, 0 to 30 mg) after therapy; 60% of patients were able to completely discontinue prednisone. Relapse occurred in 12 of the 18 patients who initially achieved remission with infliximab; 6 patients required an increase in the dose of infliximab, and steroid therapy was added in 4 patients. Adverse events that required discontinuation of infliximab therapy included abnormal liver function tests (n=1), primary histoplasmosis in a patient who traveled to an endemic region (n=1), and breast cancer (n=1).

Support for uveitis can be found In the Expert Panel Recommendation for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders. A committee of the American Uveitis Society

Reference number
1979-A

performed a systematic review of literature to generate guidelines for use of these agents in ocular inflammatory conditions. A systematic review of published studies was performed. Recommendations were generated using the Grading of Recommendations Assessment, Development, and Evaluation group criteria. Based on these studies, the expert panel recommends infliximab and adalimumab can be considered as potential second-line immunomodulatory agents for the treatment of severe ocular inflammatory conditions including posterior uveitis, panuveitis, severe uveitis associated with seronegative spondyloarthritis, and scleritis in patients requiring immunomodulation in patients who have failed or who are not candidates for antimetabolite or calcineurin inhibitor immunomodulation. Infliximab and adalimumab can be considered in these patients in preference to etanercept, which seems to be associated with lower rates of treatment success.

Support for acute graft versus host disease (GVHD) can be found in the National Comprehensive Cancer Network's guideline for hematopoietic cell transplantation. The NCCN Guideline for hematopoietic cell transplantation supports the use of infliximab in conjunction with systemic corticosteroids following no response (steroid-refractory disease) to first-line therapy options. Therapy for steroid-refractory acute GVHD is often used in conjunction with the original immunosuppressive agent.

Support for using infliximab to manage immune checkpoint inhibitor-related toxicity can be found in the National Comprehensive Cancer Network's guideline for the management of immunotherapy-related toxicities. The NCCN Guideline for the management of immunotherapy-related toxicities supports the use of adding infliximab for the management of the following immunotherapy-related conditions:

1. Myocarditis, as a further intervention if no improvement within 24 to 48 hours of starting high-dose methylprednisolone
2. Mild (G1) diarrhea or colitis if persistent or progressive symptoms and positive lactoferrin/calprotectin
3. Moderate (G2) and strongly consider for severe (G3-4) diarrhea or colitis
4. Moderate or severe inflammatory arthritis as additional disease modifying antirheumatic drug (DMARD) therapy if no improvement after holding immunotherapy and treating with oral corticosteroids or if unable to taper corticosteroids, or no response to conventional synthetic DMARDs
5. G1-4 uveitis that is refractory to high-dose systemic corticosteroids
6. Moderate (G2) pneumonitis if no improvement after 48-72 hours of corticosteroids or severe (G3-4) pneumonitis if no improvement after 48 hours of methylprednisolone
7. Stage 3 acute kidney injury/elevated serum creatinine if toxicity remains more than stage 2 after four to six weeks of corticosteroids or if creatinine increases during steroid taper (once off steroids).

Support for the use of infliximab to treat multisystem inflammatory syndrome in children (MIS-C) can be found in the COVID-19 Treatment Guidelines Panel: Coronavirus disease 2019 (COVID-19) treatment guidelines. National Institutes of Health (NIH). In pediatric patients hospitalized with multisystem inflammatory syndrome in children (MIS-C) associated with SARS-CoV-2 (COVID19), initial first line treatment is IV immune globulin with low to moderate dose glucocorticoids, such as methylprednisolone (recommendation rating, A; evidence rating, based on nonrandomized trials or observation cohort studies) and should be used in most patients (level of consensus, moderate). The risks versus benefits of treating immunocompromised MIS-C patients with immunomodulatory agents should be evaluated on an individual basis.

If MIS-C is refractory (no improvement within 24 hours of IV immune globulin and steroid initiation), initiate intensification immunomodulatory therapy (recommendation rating, A; evidence rating, expert opinion) (level of consensus, moderate) with higher-dose glucocorticoids, anakinra, or infliximab (recommendation rating, B; evidence rating, based on nonrandomized trials or observation cohort studies) (level of consensus, moderate). Infliximab should not be used in patients with MIS-C and features of macrophage activation syndrome (MAS) (level of consensus, moderate).

Severe illness may warrant dual therapy with higher-dose glucocorticoids plus anakinra (recommendation

Reference number
1979-A

rating, B; evidence rating, expert opinion), or higher-dose glucocorticoids plus infliximab (recommendation rating, B; evidence rating, expert opinion). Anakinra and infliximab should not be given in combination. Infliximab can be considered in patients with contraindications to long-term use of glucocorticoids (level of consensus, moderate). The effects of infliximab likely persist for weeks, which may provide a steroid-sparing effect.

VII. REFERENCES

1. Remicade [package insert]. Horsham, PA: Janssen Biotech, Inc.; October 2021.
2. Avsola [package insert]. Thousand Oaks, CA: Amgen; September 2021.
3. Inflectra [package insert]. New York, NY: Pfizer Inc; March 2022.
4. Renflexis [package insert]. Jersey City, NJ. Organon LLC, Inc; January 2022.
5. Infliximab [package insert]. Horsham, PA: Janssen Biotech, Inc.; October 2021.
6. van der Heijde D, Sieper J, Maksymowych WP, et al. 2010 Update of the international ASAS recommendations for the use of anti-TNF agents in patients with axial spondyloarthritis. *Ann Rheum Dis*. 2011;70:905-908.
7. IBM Micromedex® DRUGDEX® System (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at: <https://www.micromedexsolutions.com/>. Accessed August 18, 2022.
8. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol*. 2011;106(Suppl 1):S2-S25.
9. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum*. 2008;59(6):762-784.
10. Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis*. 2020;79(6):685-699. Doi:10.1136/annrheumdis-2019-216655.
11. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2016;68(1)1-26.
12. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
13. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2015 update. *Ann Rheum Dis*. 2016;75(3):499-510.
14. Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis*. 2005;64(Suppl II):ii14–ii17.
15. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol*. 2022;18(8):465-479.
16. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis*. 2011;70:896–904.
17. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care Res*. 2011;63(4):465-482.
18. Ringold S, Weiss PF, Beukelman T, et al. 2013 Update of the 2011 American College of Rheumatology Recommendations for the Treatment of Juvenile Idiopathic Arthritis: Recommendations for the Medical Therapy of Children With Systemic Juvenile Idiopathic Arthritis and Tuberculosis Screening Among Children Receiving Biologic Medications. *Arthritis & Rheumatism*. 2013;65:2499-2512.

Reference number
1979-A

19. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019;71(10):1599-1613. Doi:10.1002/art.41042.
20. The NCCN Drugs & Biologics Compendium® © 2023 National Comprehensive Cancer Network, Inc. Available at: <https://www.nccn.org>. Accessed January 16, 2023.
21. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
22. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthrit Care Res.* 2021;0:1-16.
23. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019;71(1):5-32. Doi:10.1002/art.40726.
24. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part I: Diagnosis, evaluation, and the use of complementary and procedural management. *J Am Acad Dermatol.* 2019; 81(1): 76-90.
25. Alikhan A, Sayed C, Alavi A, et al. North American clinical management guidelines for hidradenitis suppurativa: A publication from the United States and Canadian Hidradenitis Suppurativa Foundations Part II: Topical, intralesional, and systemic medical management. *J Am Acad Dermatol.* 2019; 81(1): 91-101.
26. Ringold S, Angeles-Han S, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *American College of Rheumatology.* 2019;1-18.
27. Jaffe G, Dick A, Brezin A, et al. Adalimumab in Patients with Active Noninfectious Uveitis. *N Engl J Med.* 2016; 375: 932-943.
28. Nguyen QD, Merrill P, Jaffe G, et al. Adalimumab for prevention of uveitic flare in patients with inactive non-infectious uveitis controlled by corticosteroids (VISUAL II): a multicentre, double-masked, randomized, placebo-controlled phase 3 trial. *Lancet.* 2016; 17: 388(10050):1183-92.
29. Feuerstein J, Ho E, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology.* 2021; 160:2496-2508.
30. Elmets C, Korman N, et al. Guidelines of Care for the Management and Treatment of Psoriasis with Topical Therapy and Alternative Medicine Modalities for Psoriasis Severity Measures. *J Am Acad Dermatol.* 2021; 84:432-470.
31. Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* 2018;113:481-517.
32. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology.* 2020; 158:1450.
33. Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019;114:384-413.
34. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
35. Elmets C, Korman N, et al. Guidelines of Care for the Management and Treatment of Psoriasis with Topical Therapy and Alternative Medicine Modalities for Psoriasis Severity Measures. *J Am Acad Dermatol.* 2021;84:432-470.

Reference number
1979-A

36. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol.* 2022;74(4):553-569.
37. Mehta NA, Emani-Naeini P. A review of systemic biologics and local immunosuppressive medications in uveitis. *J Ophthalmic Vis Res.* 2022;17(2):276-289.
38. Hatemi G et al. EULAR recommendations on management of Behcet syndrome. *Ann Rheum Dis* 2018 Jun;77(6):808
39. Kraetsch HG, Antoni C, Kalden JR, et al: Successful treatment of a small cohort of patients with adult onset of Still's disease with infliximab: first experiences. *Ann Rheum Dis* 2001; 60:iii55-iii57.
40. Cavagna L, Caporali R, Epis O, et al: Infliximab in the treatment of adult Still's disease refractory to conventional therapy. *Clin Exp Rheumatol* 2001; 19:329-332.
41. Ellman MH, Hanauer S, Sitrin M, et al: Crohn's disease arthritis treated with infliximab: an open trial in four patients. *J Clin Rheumatol* 2001; 7:67-71.
42. Ingram JR, Collier F, Brown D, et al. British Association of Dermatologists guidelines for the management of hidradenitis suppurativa (acne inversa) 2018. *Br J Dermatol.* 2019 May;180(5):1009-1017
43. Pascher A, Padke C, Dignass A, et al: Successful infliximab treatment of steroid and OKT3 refractory acute cellular rejection in two patients after intestinal transplantation. *Transplantation* 2003; 76(3):615-618.
44. Cantini F, Niccoli L, Salvarani C, et al: Treatment of longstanding active giant cell arteritis with infliximab: a report of 4 cases. *Arthritis Rheum* 2001; 44(12):2933-2935.
45. McCrindle BW, Rowley AH, Newburger JW, et al: Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals from the American Heart Association. *Circulation* 2017; 135(17):e927-e999.
46. Kolde G, Muche JM, Schulze P, et al: Infliximab: a promising new treatment option for ulcerated necrobiosis lipoidica. *Dermatology* 2003; 206:180-181.
47. Ahronowitz I, Harp J, Shinkai K. Etiology and management of pyoderma gangrenosum: a comprehensive review. *Am J Clin Dermatol.* 2012 Jun 1;13(3):191-211
48. Baughman RP, Valeyre D, Korsten P, et al: ERS clinical practice guidelines on treatment of sarcoidosis. *Eur Respir J* 2021.
49. Voigtlander C, Luftl M, Schuler G, et al: Infliximab (anti-tumor necrosis factor alpha antibody): a novel, highly effective treatment of recalcitrant subcorneal pustular dermatosis (Sneddon-Wilkinson Disease). *Arch Dermatol* 2001; 137:1571-1574.
50. Taylor PC, Steuer A, Gruber J, et al: Comparison of ultrasonographic assessment of synovitis and joint vascularity with radiographic evaluation in a randomized, placebo-controlled study of infliximab therapy in early rheumatoid arthritis. *Arthritis Rheum* 2004; 50(4):1107-1116.
51. Quinn MA , Conaghan PG , O'Connor PJ , et al: Very early treatment with infliximab in addition to methotrexate in early, poor-prognosis rheumatoid arthritis reduces magnetic resonance imaging evidence of synovitis and damage, with sustained benefit after infliximab withdrawal: results from a twelve-month randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2005; 52(1):27-35.
52. Pipitone N, Olivieri I, & Salvarani C: Recommendations of the Italian Society of Rheumatology for the treatment of the primary large-vessel vasculitis with biological agents. *Clin Exp Rheumatol* 2012; 30(SUPPL. 70):S139-S161.
53. Mekinian A, Neel A, Sibilia J, et al: Efficacy and tolerance of infliximab in refractory Takayasu arteritis: French multicentre study. *Rheumatology (Oxford)* 2012; 51(5):882-886.
54. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders. *Ophthalmology.* 2014;121(3):785-796.
55. COVID-19 Treatment Guidelines Panel: Coronavirus disease 2019 (COVID-19) treatment guidelines. National Institutes of Health (NIH). Bethesda, MD. 2022. Accessed July 3, 2023.