

STANDARD MEDICARE PART B MANAGEMENT

MIRCERA (methoxy polyethylene glycol-epoetin beta)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Mircera is indicated for the treatment of anemia associated with chronic kidney disease (CKD) in:

- Adult patients on dialysis and adult patients not on dialysis.
- Pediatric patients 5 to 17 years of age on hemodialysis who are converting from another erythropoiesis-stimulating agent (ESA) after their hemoglobin level was stabilized with an ESA.

Limitations of Use:

Mircera is not indicated and is not recommended:

- In the treatment of anemia due to cancer chemotherapy
- As a substitute for red blood cell (RBC) transfusions in patients who require immediate correction of anemia

Mircera has not been shown to improve symptoms, physical functioning, or health-related quality of life.

Note: Use in members on dialysis is covered under the Medicare Part B dialysis benefit and is excluded from coverage under this policy

All other indications will be assessed on an individual basis. Submissions for indications other than those enumerated in this policy should be accompanied by supporting evidence from Medicare approved compendia.

II. CRITERIA FOR INITIAL APPROVAL

Note: Requirements regarding hemoglobin level exclude values due to recent transfusion.

Anemia Due to Chronic Kidney Disease (CKD)

Authorization of 12 weeks may be granted for the treatment of anemia due to chronic kidney disease in members not receiving dialysis with a pretreatment hemoglobin of less than 10 g/dL or a hematocrit of less than 30%.

III. CONTINUATION OF THERAPY

Note: Requirements regarding current hemoglobin level exclude values due to recent transfusion.

Reference number(s)
4462-A

All members (including new members) requesting authorization for continuation of therapy must be currently receiving therapy with the requested agent.

Authorization for 12 weeks may be granted when all of the following criteria are met:

- A. The member is currently receiving therapy with Mircera.
- B. Mircera is being used to treat anemia due to chronic kidney disease (CKD).
- C. Mircera has been effective for treating the diagnosis or condition.

IV. SUMMARY OF EVIDENCE

The contents of this policy were created after examining the following resources:

- 1. The prescribing information for Mircera.
- 2. The available compendium
 - a. National Comprehensive Cancer Network (NCCN) Drugs and Biologics Compendium
 - b. Micromedex DrugDex
 - c. American Hospital Formulary Service- Drug Information (AHFS-DI)
 - d. Lexi-Drugs
 - e. Clinical Pharmacology
- 3. Kidney Disease: Improving Global Outcomes (KDIGO) 2012 Clinical Practice Guideline for Anemia for Chronic Kidney Disease

After reviewing the information in the above resources, the FDA-approved indications listed in the prescribing information for Mircera are covered.

V. EXPLANATION OF RATIONALE

Support for FDA-approved indications can be found in the manufacturer's prescribing information.

VI. REFERENCES

- 1. Mircera [package insert]. St. Gallen, Switzerland: Vifor (International) Inc.; March 2023.
- 2. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Int.* 2012;Suppl 2:279-335.