

Reference number
2051-A

# SPECIALTY GUIDELINE MANAGEMENT

## CEREZYME (imiglucerase)

### POLICY

#### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

Cerezyme is indicated for treatment of adults and pediatric patients 2 years of age and older with Type 1 Gaucher disease that results in one or more of the following conditions: anemia, thrombocytopenia, bone disease, and/or hepatomegaly or splenomegaly.

B. Compendial Uses

1. Gaucher disease type 2
2. Gaucher disease type 3

All other indications are considered experimental/investigational and not medically necessary.

#### II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review: beta-glucocerebrosidase enzyme assay or genetic testing results supporting diagnosis.

#### III. CRITERIA FOR INITIAL APPROVAL

A. **Gaucher disease type 1**

Authorization of 12 months may be granted for treatment of Gaucher disease type 1 when the diagnosis of Gaucher disease was confirmed by enzyme assay demonstrating a deficiency of beta-glucocerebrosidase (glucosidase) enzyme activity or by genetic testing.

B. **Gaucher disease type 2**

Authorization of 12 months may be granted for treatment of Gaucher disease type 2 when the diagnosis of Gaucher disease was confirmed by enzyme assay demonstrating a deficiency of beta-glucocerebrosidase (glucosidase) enzyme activity or by genetic testing.

C. **Gaucher disease type 3**

Authorization of 12 months may be granted for treatment of Gaucher disease type 3 when the diagnosis of Gaucher disease was confirmed by enzyme assay demonstrating a deficiency of beta-glucocerebrosidase (glucosidase) enzyme activity or by genetic testing.

#### IV. CONTINUATION OF THERAPY

<b>Reference number</b>
2051-A

Authorization of 12 months may be granted for continued treatment of an indication listed in Section III when all of the following criteria are met:

- A. Member meets the criteria for initial approval.
- B. Member is not experiencing an inadequate response or any intolerable adverse events from therapy.

## V. REFERENCES

1. Cerezyme [package insert]. Cambridge, MA: Genzyme Corporation; December 2021.
2. Altarescu G, Hill S, Wiggs E, et al. The efficacy of enzyme replacement therapy in patients with chronic neuronopathic Gaucher's disease. *J Pediatr*. 2001;138:539-547.
3. Erikson A, Forsberg H, Nilsson M, Astrom M, Mansson JE. Ten years' experience of enzyme infusion therapy of Norrbottnian (type 3) Gaucher disease. *Acta Paediatr*. 2006;95:312-317.
4. Pastores GM, Hughes DA. Gaucher Disease. 2000 July 27 [Updated June 21, 2018]. In: Adam MP, Everman DB, Mirzaa GM, et al, editors. GeneReviews® [Internet]. Seattle, WA: University of Washington, Seattle; 1993-2022.
5. Kaplan P, Baris H, De Meirleir L, et al. Revised recommendations for the management of Gaucher disease in children. *Eur J Pediatr*. 2013;172:447-458.
6. National Organization for Rare Disorders. (2003). *NORD guide to rare disorders*. Philadelphia: Lippincott Williams & Wilkins.