

SPECIALTY GUIDELINE MANAGEMENT

ENTYVIO (vedolizumab)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Adult patients with moderately to severely active ulcerative colitis (UC).
2. Adult patients with moderately to severely active Crohn's disease (CD).

B. Compendial Uses

Immune checkpoint inhibitor-related toxicity

All other indications are considered experimental/investigational and not medically necessary.

II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

A. Ulcerative colitis (UC) and Crohn's disease (CD)

Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.

B. Immune checkpoint inhibitor-related toxicity

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

III. PRESCRIBER SPECIALTIES

The medication must be prescribed by or in consultation with one of the following:

- A. Crohn's disease and ulcerative colitis: gastroenterologist
- B. Immune checkpoint inhibitor-related toxicity: gastroenterologist, hematologist, or oncologist

IV. CRITERIA FOR INITIAL APPROVAL

A. **Ulcerative colitis (UC)**

Authorization of 12 months may be granted for adult members for treatment of moderately to severely active ulcerative colitis.

B. **Crohn's disease (CD)**

Authorization of 12 months may be granted for adult members for treatment of moderately to severely active Crohn's disease.

C. Immune checkpoint inhibitor-related toxicity

Authorization of 6 months may be granted for the treatment of immune checkpoint inhibitor-related diarrhea or colitis when the member has experienced an inadequate response, intolerance, or contraindication to systemic corticosteroids or infliximab.

V. CONTINUATION OF THERAPY

A. Ulcerative colitis (UC)

1. Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Stool frequency
 - ii. Rectal bleeding
 - iii. Urgency of defecation
 - iv. C-reactive protein (CRP)
 - v. Fecal calprotectin (FC)
 - vi. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
 - vii. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

B. Crohn's disease (CD)

1. Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active Crohn's disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Abdominal pain or tenderness
 - ii. Diarrhea
 - iii. Body weight
 - iv. Abdominal mass
 - v. Hematocrit
 - vi. Appearance of the mucosa on endoscopy, computed tomography enterography (CTE), magnetic resonance enterography (MRE), or intestinal ultrasound
 - vii. Improvement on a disease activity scoring tool (e.g., Crohn's Disease Activity Index [CDAI] score)

C. Immune checkpoint inhibitor-related toxicity

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

Reference number(s)
2004-A

VI. OTHER

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

VII. DOSAGE AND ADMINISTRATION

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

VIII. REFERENCES

1. Entyvio [package insert]. Lexington, MA: Takeda Pharmaceuticals U.S.A., Inc.; September 2023.
2. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol.* 2011;106(Suppl 1):S2-S25.
3. Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* 2018;113:481-517.
4. Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol.* 2019;114:384-413.
5. The NCCN Drugs & Biologics Compendium® © 2023 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed January 11, 2024.
6. NCCN Clinical Practice Guidelines in Oncology® (NCCN Guidelines®). Management of Immunotherapy-Related Toxicities. Version 1.2024. Available at: www.nccn.org. Accessed January 11, 2024.
7. Schneider BJ, Naidoo J, Santomaso BD, et al. Management of Immune-Related Adverse Events in Patients Treated With Immune Checkpoint Inhibitor Therapy: American Society of Clinical Oncology Guideline Update. *J Clin Oncol.* 2021; 39(36):4073-4126.
8. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology.* 2020; 158:1450-1461.
9. Feuerstein JD, Ho EY, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology.* 2021; 160: 2496-2508.
10. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-1486.