

Reference number(s)
1894-A

# SPECIALTY GUIDELINE MANAGEMENT

## OPDIVO (nivolumab)

### POLICY

#### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### A. FDA-Approved Indications

1. **Unresectable or Metastatic Melanoma**  
Opdivo, as a single agent or in combination with ipilimumab, is indicated for the treatment of adult and pediatric patients 12 years and older with unresectable or metastatic melanoma.
2. **Adjuvant Treatment of Melanoma**  
Opdivo is indicated for the adjuvant treatment of adult and pediatric patients 12 years and older with completely resected stage IIB, stage IIC, stage III, or stage IV melanoma.
3. **Metastatic Non-Small Cell Lung Cancer**
  - a. Opdivo, in combination with ipilimumab, is indicated for the first-line treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors express PD-L1 ( $\geq 1\%$ ) as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations.
  - b. Opdivo, in combination with ipilimumab and 2 cycles of platinum-doublet chemotherapy, is indicated for the first-line treatment of adult patients with metastatic or recurrent NSCLC, with no EGFR or ALK genomic tumor aberrations.
  - c. Opdivo is indicated for the treatment of adult patients with metastatic NSCLC with progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving Opdivo.
4. **Neoadjuvant Treatment of Resectable Non-Small Cell Lung Cancer**  
Opdivo, in combination with platinum-doublet chemotherapy, is indicated as neoadjuvant treatment of adult patients with resectable (tumors  $\geq 4$  cm or node positive) non-small cell lung cancer (NSCLC).
5. **Malignant Pleural Mesothelioma**  
Opdivo, in combination with ipilimumab, is indicated for the first-line treatment of adult patients with unresectable malignant pleural mesothelioma.
6. **Advanced Renal Cell Carcinoma**
  - a. Opdivo as a single agent is indicated for the treatment of adult patients with advanced renal cell carcinoma (RCC) who have received prior anti-angiogenic therapy.
  - b. Opdivo, in combination with ipilimumab, is indicated for the first-line treatment of adult patients with intermediate or poor risk advanced RCC.
  - c. Opdivo, in combination with cabozantinib, is indicated for the first-line treatment of adult patients with advanced RCC.
7. **Classical Hodgkin Lymphoma**

Reference number(s)
1894-A

Opdivo is indicated for the treatment of adult patients with classical Hodgkin lymphoma (cHL) that has relapsed or progressed after:

- a. Autologous hematopoietic stem cell transplantation (HSCT) and brentuximab vedotin, or
- b. Three or more lines of systemic therapy that includes autologous HSCT.

8. Squamous Cell Carcinoma of the Head and Neck

Opdivo is indicated for the treatment of adult patients with recurrent or metastatic squamous cell carcinoma of the head and neck (SCCHN) with disease progression on or after platinum-based therapy.

9. Urothelial Carcinoma

- a. Opdivo is indicated for the adjuvant treatment of adult patients with urothelial carcinoma (UC) who are at high risk of recurrence after undergoing radical resection of UC.
- b. Opdivo, in combination with cisplatin and gemcitabine, is indicated for the first-line treatment of adult patients with unresectable or metastatic urothelial carcinoma.
- c. Opdivo is indicated for the treatment of adult patients with locally advanced or metastatic urothelial carcinoma who:
  - i. Have disease progression during or following platinum-containing chemotherapy
  - ii. Have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy.

10. Microsatellite Instability-High or Mismatch Repair Deficient Metastatic Colorectal Cancer

Opdivo, as a single agent or in combination with ipilimumab, is indicated for the treatment of adult and pediatric patients 12 years and older with microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) metastatic colorectal cancer (CRC) that has progressed following treatment with a fluoropyrimidine, oxaliplatin, and irinotecan.

11. Hepatocellular Carcinoma

Opdivo, in combination with ipilimumab, is indicated for the treatment of adult patients with hepatocellular carcinoma (HCC) who have been previously treated with sorafenib.

12. Esophageal Carcinoma

- a. Opdivo is indicated for the adjuvant treatment of completely resected esophageal or gastroesophageal junction cancer with residual pathologic disease in adult patients who have received neoadjuvant chemoradiotherapy (CRT).
- b. Opdivo, in combination with fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the first-line treatment of adult patients with unresectable advanced or metastatic esophageal squamous cell carcinoma (ESCC).
- c. Opdivo, in combination with ipilimumab, is indicated for the first-line treatment of adult patients with unresectable advanced or metastatic esophageal squamous cell carcinoma (ESCC).
- d. Opdivo is indicated for the treatment of adult patients with unresectable advanced, recurrent or metastatic esophageal squamous cell carcinoma (ESCC) after prior fluoropyrimidine- and platinum-based chemotherapy.

13. Gastric Cancer, Gastroesophageal Junction Cancer, Esophageal Adenocarcinoma

Opdivo, in combination with fluoropyrimidine- and platinum-containing chemotherapy, is indicated for the treatment of adult patients with advanced or metastatic gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma.

**B. Compendial Uses**

1. Cutaneous melanoma
2. Non-small cell lung cancer

Reference number(s)
1894-A

3. Renal cell carcinoma
4. Classical Hodgkin lymphoma
5. Head and neck cancers
6. Urothelial carcinoma
  - a. Bladder cancer
  - b. Primary carcinoma of the urethra
  - c. Upper genitourinary tract tumors
  - d. Urothelial carcinoma of the prostate
7. Colorectal cancer, including appendiceal adenocarcinoma and anal adenocarcinoma
8. Hepatocellular carcinoma
9. Uveal Melanoma
10. Anal Carcinoma
11. Merkel Cell Carcinoma
12. Central Nervous System (CNS) brain metastases
13. Gestational trophoblastic neoplasia
14. Malignant pleural mesothelioma
15. Malignant peritoneal mesothelioma
16. Small bowel adenocarcinoma
17. Ampullary Adenocarcinoma
18. Extranodal NK/T-cell lymphoma
19. Endometrial Carcinoma
20. Vulvar Cancer
21. Gastric Cancer
22. Esophageal/Esophagogastric Junction Cancers
23. Small cell lung cancer
24. Cervical Cancer
25. Pediatric Diffuse High-Grade Gliomas
26. Pediatric Primary Mediastinal Large B-cell Lymphoma
27. Kaposi Sarcoma
28. Bone Cancer
29. Biliary Tract Cancers
  - a. Cholangiocarcinoma
  - b. Gallbladder Cancer
30. Soft Tissue Sarcoma
  - a. Extremity/body wall sarcoma
  - b. Head/neck sarcoma
  - c. Retroperitoneal/intra-abdominal sarcoma
  - d. Rhabdomyosarcoma
  - e. Angiosarcoma

All other indications are considered experimental/investigational and not medically necessary.

## II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

- A. Documentation of laboratory report confirming MSI-H or mismatch repair deficient (dMMR) tumor status, where applicable.
- B. Documentation of programmed death ligand 1 (PD-L1) tumor expression, where applicable.
- C. Documentation of the presence of EGFR exon 19 deletions or exon 21 L858R mutations or ALK rearrangements, where applicable.

### III. EXCLUSIONS

Coverage will not be provided for members who have experienced disease progression while on programmed death receptor-1 (PD-1) or programmed death ligand 1 (PD-L1) inhibitor therapy (other than when used as second-line or subsequent therapy for metastatic or unresectable melanoma in combination with ipilimumab following progression on single agent anti-PD-1 immunotherapy).

### IV. CRITERIA FOR INITIAL APPROVAL

#### A. Cutaneous Melanoma

Authorization of 6 months may be granted for treatment of cutaneous melanoma in either of the following settings:

1. The requested medication will be used as a single agent or in combination with ipilimumab (4 doses of ipilimumab, followed by Opdivo as a single agent) for locally recurrent, unresectable, progressive or metastatic disease.
2. The requested medication will be used as a single agent or in combination with ipilimumab (4 doses of ipilimumab, followed by Opdivo as a single agent) as adjuvant treatment of stage III or IV disease following complete resection or no evidence of disease.
3. The requested medication will be used as a single agent as adjuvant treatment of stage IIB and IIC disease following complete resection.

#### B. Non-Small Cell Lung Cancer (NSCLC)

1. Authorization of 6 months may be granted for treatment of recurrent, advanced or metastatic non-small cell lung cancer if either of the following criteria are met:
  - a. There are no EGFR exon 19 deletions or exon 21 L858R mutations or ALK rearrangements (unless testing is not feasible due to insufficient tissue) and the requested medication will be used in a regimen containing ipilimumab.
  - b. The requested medication will be used as single agent subsequent therapy.
2. Authorization of 3 months (for up to 3 cycles total) may be granted for neoadjuvant treatment of resectable non-small cell lung cancer (NSCLC) in combination with platinum-doublet chemotherapy.

#### C. Renal Cell Carcinoma

Authorization of 6 months may be granted for treatment of relapsed, advanced, or stage IV renal cell carcinoma, in any of the following settings:

1. The requested medication will be used as a single agent for clear cell histology as subsequent therapy.
2. The requested medication will be used as a single agent for non-clear cell histology.
3. The requested medication will be used in combination with ipilimumab (4 doses of ipilimumab, followed by Opdivo as a single agent) for disease with clear cell histology as:
  - a. First-line therapy for poor or intermediate risk.
  - b. First-line therapy for favorable risk.
  - c. Subsequent therapy.
4. The requested medication will be used in combination with cabozantinib.

#### D. Classical Hodgkin Lymphoma (cHL)

Authorization of 6 months may be granted for treatment of classical Hodgkin lymphoma when either of the following criteria is met:

1. The requested medication will be used as palliative or subsequent therapy and the member meets any of the following criteria:
  - a. Member has relapsed or progressed after high-dose therapy and autologous stem cell rescue (HDT/ASCR).

Reference number(s)
1894-A

- b. Member has relapsed or refractory disease and is transplant-ineligible.
- c. Member has relapsed or refractory disease and was either heavily pretreated or there was a decrease in cardiac function.
- d. Member is post-allogeneic transplant
2. The requested medication will be used in combination with brentuximab vedotin or in combination with ICE (ifosfamide, carboplatin, etoposide) for relapsed or refractory disease.
3. The requested medication will be used as a single agent for disease refractory to at least three lines of prior therapy.

#### **E. Head and Neck Cancers**

Authorization of 6 months may be granted for treatment of head and neck cancers in members who meet either of the following criteria:

1. For unresectable, recurrent, persistent or metastatic disease.
2. For nasopharyngeal cancer in combination with cisplatin and gemcitabine for unresectable, recurrent, persistent or metastatic disease.

#### **F. Urothelial Carcinoma – Bladder Cancer**

1. Authorization of 6 months may be granted in combination with gemcitabine and cisplatin for up to 6 cycles followed by nivolumab maintenance therapy as first line treatment of bladder cancer.
2. Authorization of 6 months may be granted as a single agent for treatment of bladder cancer when any of the following conditions are met:
  - a. As subsequent therapy for locally advanced, recurrent, persistent, or metastatic disease.
  - b. As adjuvant therapy in members who are at high risk of recurrence after undergoing resection.

#### **G. Urothelial Carcinoma – Primary Carcinoma of the Urethra**

1. Authorization of 6 months may be granted in combination with gemcitabine and cisplatin for up to 6 cycles followed by nivolumab maintenance therapy as first line treatment of primary carcinoma of the urethra.
2. Authorization of 6 months may be granted as a single agent for treatment of primary carcinoma of the urethra when either of the following are met:
  - a. As subsequent therapy for recurrent, locally advanced, or metastatic disease.
  - b. As adjuvant therapy in members who are at high risk of recurrence after undergoing resection.

#### **H. Urothelial Carcinoma – Upper Genitourinary Tract Tumors or Urothelial Carcinoma of the Prostate**

1. Authorization of 6 months may be granted in combination with gemcitabine and cisplatin for up to 6 cycles followed by nivolumab maintenance therapy as first line treatment of metastatic upper genitourinary (GU) tract tumors or urothelial carcinoma of the prostate.
2. Authorization of 6 months may be granted as a single agent for treatment of upper genitourinary (GU) tract tumors or urothelial carcinoma of the prostate when either of the following are met:
  - a. As subsequent therapy for locally advanced or metastatic disease.
  - b. As adjuvant therapy in members who are at high risk of recurrence after undergoing resection.

#### **I. Colorectal Cancer**

Authorization of 6 months may be granted for treatment of colorectal cancer, including appendiceal adenocarcinoma and anal adenocarcinoma, for microsatellite-instability high (MSI-H) or mismatch repair deficient (dMMR) tumors when used as a single agent or in combination with ipilimumab (4 doses of ipilimumab, followed by Opdivo as a single agent) for advanced, metastatic, unresectable, or inoperable disease.

#### **J. Small Bowel Adenocarcinoma**

Reference number(s)
1894-A

Authorization of 6 months may be granted as a single agent or in combination with ipilimumab for treatment of advanced or metastatic small bowel adenocarcinoma for microsatellite-instability high (MSI-H) or mismatch repair deficient (dMMR) tumors.

**K. Ampullary Adenocarcinoma**

Authorization of 6 months may be granted in combination with ipilimumab for treatment of progressive, unresectable, or metastatic microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) ampullary adenocarcinoma.

**L. Hepatocellular Carcinoma**

Authorization of 6 months may be granted as a single agent or in combination with ipilimumab (4 doses of ipilimumab, followed by Opdivo as a single agent) for treatment of hepatocellular carcinoma.

**M. Uveal Melanoma**

Authorization of 6 months may be granted as a single agent or in combination with ipilimumab for treatment of uveal melanoma for unresectable or metastatic disease.

**N. Anal Carcinoma**

Authorization of 6 months may be granted as a single agent for subsequent treatment of metastatic anal carcinoma.

**O. Merkel Cell Carcinoma**

Authorization of 6 months may be granted for treatment of Merkel cell carcinoma in either of the following settings:

1. Metastatic disease.
2. Neoadjuvant treatment of node positive disease.
3. Progressive, unresectable, recurrent, or stage IV disease when used in combination with ipilimumab.

**P. CNS Brain Metastases**

Authorization of 6 months may be granted for treatment of CNS brain metastases when either of the following criteria are met:

1. The requested medication will be used as a single agent or in combination with ipilimumab in members with melanoma.
2. The requested medication will be used as a single agent in members with PD-L1 positive non-small cell lung cancer.

**Q. Gestational Trophoblastic Neoplasia**

Authorization of 6 months may be granted as a single agent for treatment of gestational trophoblastic neoplasia for multiagent chemotherapy-resistant disease when either of the following criteria is met:

1. Member has recurrent or progressive intermediate trophoblastic tumor (placental site trophoblastic tumor or epithelioid trophoblastic tumor) following treatment with a platinum-based regimen.
2. Member has high-risk disease.

**R. Malignant Pleural or Peritoneal Mesothelioma**

Authorization of 6 months may be granted for the treatment of malignant pleural or peritoneal mesothelioma, including pericardial mesothelioma and tunica vaginalis testis mesothelioma, in either of the following settings:

1. The requested medication will be used as first line therapy in combination with ipilimumab.
2. The requested medication will be used as subsequent therapy as a single agent or in combination with ipilimumab.

**S. Esophageal and Esophagogastric Junction Carcinoma**



1. Authorization of 6 months may be granted for treatment of esophageal or esophagogastric junction in members who are not surgical candidates or have unresectable locally advanced, recurrent or metastatic disease when the requested medication will be used in combination with ipilimumab or fluoropyrimidine and platinum containing chemotherapy or will be used as subsequent therapy.
2. Authorization of 6 months may be granted for adjuvant treatment of completely resected esophageal or esophagogastric junction cancer with residual pathologic disease.
3. Authorization of 6 months may be granted as a single agent or in combination with ipilimumab for neoadjuvant or perioperative treatment of esophageal or esophagogastric junction adenocarcinoma if tumor is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) and member is medically fit for surgery.

**T. Extranodal NK/T-Cell Lymphoma**

Authorization of 6 months may be granted for treatment of relapsed or refractory extranodal NK/T-cell lymphoma.

**U. Endometrial Carcinoma**

Authorization of 6 months may be granted as a single agent for subsequent treatment of recurrent or metastatic microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) endometrial carcinoma.

**V. Vulvar Cancer**

Authorization of 6 months may be granted for treatment of HPV-related advanced, recurrent, or metastatic vulvar cancer as subsequent therapy as a single agent.

**W. Gastric Cancer**

Authorization of 6 months may be granted for treatment of gastric cancer in any of the following settings:

1. When the requested medication is being used in members who are not surgical candidates or have unresectable locally advanced, recurrent, or metastatic disease, when the requested medication will be used in combination with ipilimumab or chemotherapy.
2. When the requested medication will be used as a single agent or in combination with ipilimumab for neoadjuvant or perioperative treatment of gastric adenocarcinoma if tumor is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) and member is medically fit for surgery

**X. Small Cell Lung Cancer**

Authorization of 6 months may be granted for subsequent treatment of relapsed or progressive small cell lung cancer as a single agent.

**Y. Cervical Cancer**

Authorization of 6 months may be granted for subsequent treatment of recurrent or metastatic cervical cancer as a single agent if PD-L1 positive (combined positive score [CPS] ≥1).

**Z. Pediatric Diffuse High-Grade Gliomas**

Authorization of 6 months may be granted for hypermutant tumor pediatric diffuse high-grade glioma as adjuvant treatment or for recurrent or progressive disease.

**AA. Pediatric Primary Mediastinal Large B-Cell Lymphoma**

Authorization of 6 months may be granted as a single agent or in combination with brentuximab vedotin for treatment of relapsed or refractory primary mediastinal large B-cell lymphoma.

**BB. Kaposi Sarcoma**

Authorization of 6 months may be granted in combination with ipilimumab for subsequent treatment of relapsed/refractory classic Kaposi Sarcoma.

Reference number(s)
1894-A

### **CC. Bone Cancer**

Authorization of 6 months may be granted in combination with ipilimumab for unresectable or metastatic disease when all of the following are met:

1. Disease has tumor mutation burden-high (TMB-H) [ $\geq 10$  mutations/megabase (mut/Mb)] tumors
2. Disease has progressed following prior treatment and has no satisfactory alternative treatment options

### **DD. Biliary Tract Cancers (Cholangiocarcinoma and Gallbladder Cancer)**

Authorization of 6 months may be granted as subsequent treatment in combination with ipilimumab for unresectable or resected gross residual (R2) disease, or metastatic disease that is tumor mutation burden-high (TMB-H).

### **EE. Soft Tissue Sarcoma**

Authorization of 6 months may be granted for treatment of soft tissue sarcoma in the following settings:

1. The requested medication will be used as a single agent or in combination with ipilimumab for treatment of extremity/body wall sarcomas, head/neck sarcomas and retroperitoneal/intra-abdominal sarcomas and rhabdomyosarcoma.
2. The requested medication will be used in combination with ipilimumab for the treatment of angiosarcoma.

## **V. CONTINUATION OF THERAPY**

### **A. Adjuvant treatment of melanoma**

Authorization of 6 months may be granted (up to 12 months total) for continued treatment in members requesting reauthorization for cutaneous melanoma who have not experienced disease recurrence or an unacceptable toxicity.

### **B. Urothelial carcinoma**

1. Authorization of 6 months may be granted (up to 12 months total) for continued treatment in members requesting reauthorization for adjuvant treatment of urothelial carcinoma who have not experienced disease recurrence or an unacceptable toxicity.
2. Authorization of 6 months may be granted (up to 24 months total) for continued treatment in members requesting reauthorization for urothelial carcinoma when the requested medication is used in combination with gemcitabine and cisplatin for up to 6 cycles followed by nivolumab maintenance therapy when the member has not experienced disease progression or an unacceptable toxicity.

### **C. Non-small cell lung cancer or Malignant pleural mesothelioma**

Authorization of 6 months may be granted (up to 24 months total when used in combination with ipilimumab) for continued treatment in members requesting reauthorization for non-small cell lung cancer or malignant pleural, including pericardial mesothelioma and tunica vaginalis testis mesothelioma subtypes, when there is no evidence of unacceptable toxicity or disease progression while on the current regimen. Neoadjuvant treatment of NSCLC will be approved for a total of 3 months of therapy.

### **D. Renal Cell Carcinoma**

Authorization of 6 months may be granted (up to 24 months total when used in combination with cabozantinib) for continued treatment in members requesting reauthorization for renal cell carcinoma when there is no evidence of unacceptable toxicity or disease progression while on the current regimen.

### **E. Gastric Cancer, Esophageal Cancer, and Esophagogastric Junction Carcinoma**



Reference number(s)
1894-A

Authorization of 6 months may be granted for continued treatment in members requesting reauthorization for gastric cancer, esophageal cancer, and esophagogastric junction carcinoma when there is no evidence of unacceptable toxicity or disease progression while on the current regimen for the following durations of therapy:

1. Esophageal squamous cell carcinoma in combination with ipilimumab or chemotherapy for up to 24 months
2. Unresectable advanced, recurrent or metastatic esophageal squamous cell carcinoma as a single agent until disease progression or unacceptable toxicity
3. Adjuvant treatment of resected esophageal or esophagogastric junction cancer as a single agent for up to 12 months
4. Gastric cancer, esophagogastric junction cancer, and esophageal adenocarcinoma in combination with chemotherapy for up to 24 months

#### **F. All other indications**

Authorization of 6 months may be granted for continued treatment in members requesting reauthorization for all other indications listed in Section IV when there is no evidence of unacceptable toxicity or disease progression while on the current regimen.

#### **VI. REFERENCES**

1. Opdivo [package insert]. Princeton, NJ: Bristol-Myers Squibb Company; March 2024.
2. The NCCN Drugs & Biologics Compendium® © 2024 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed March 4, 2024.
3. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Anal Carcinoma. Version 3.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/anal.pdf](https://www.nccn.org/professionals/physician_gls/pdf/anal.pdf) Accessed September 6, 2023
4. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Melanoma: Cutaneous. Version 2.2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/cutaneous\\_melanoma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cutaneous_melanoma.pdf). Accessed September 6, 2023
5. Lexicomp [database online]. Hudson, OH: Lexi-Comp, Inc.; <https://online.lexi.com/lco/action/home> [available with subscription]. Accessed September 6, 2023.