

Preventing Teen Pregnancy Infographics

Vital Signs

Updated Apr. 7, 2015



Bar chart shows how use of long-acting reversible contraception (LARC) among teens ages 15-19 has increased in total from less than 1% in 2005 to 7% in 2013 at Title X-funded centers. It also show preference between the two LARC methods intrauterine device (IUD) and implant.



How does use of LARC vary by state among teens ages 15-19 seeking birth control at Title X-funded centers?

0-5%

- Alabama
- Arkansas
- Delaware
- Florida
- Georgia
- 6-10% • Arizona
- California
- Connecticut
- Illinois
- Maine

- 11-20%
- Alaska
- District of Columbia
- Hawaii
- Iowa
- New Hampshire

- 21% +
 - Colorado

https://archive.cdc.gov/#/details?url=https://www.cdc.gov/vitalsigns/larc/infographic.html

- Idaho
- Indiana
- Kansas
- Kentucky
- Louisiana
- Michigan
- Mississippi
- Missouri
- Montana
- Nevada
- New Jersey
- North Dakota
- Ohio
- Pennsylvania
- South Dakota
- Utah
- West Virginia
- Wyoming

- Oregon
 - Rhode Island
- Vermont
 - Washington
- New Mexico •

Nebraska

Minnesota

• Maryland

• Massachusetts

New York •

•

- North Carolina •
 - Oklahoma ٠
 - South Carolina •
 - Tennessee ٠
 - Texas
 - Virginia
 - Wisconsin

- How effective is LARC at preventing pregnancy compared with other birth control commonly used by teens?
- IMPLANT IUD **MALE CONDOM** PILL HOW TO USE IT: HOW TO USE IT: HOW TO USE IT: HOW TO USE IT: · Use correctly every · Placed by health care provider · Placed by health care provider Take at the same time time during sex each day · Lasts up to 3 years · Copper IUD lasts up to 10 years Progestin IUD lasts 3–5 years Chances of getting pregnant: Chances of getting pregnant: Chances of getting pregnant: Less than 1 out of 100 women* Less than 1 out of 100 women

tess than 1 out of 100 women	Less than 1 out of 100 women	9 out of 100 women*	18 out of 100 women*
	TIVE		LEAST EFFECTIVE
	oms should always be u ntrol to protect against	-	
*Number of pregnancies per 10	0 women using the method within first	year of typical use.	
URCE: Trussell J., Contraception, M	May 2011; www.cdc.gov/reproductivehea	alth/UnintendedPregnancy/Contracept	ion htm

How effective is LARC at preventing pregnancy compared with other birth control commonly used by teens?

(In order from Most Effective to Least Effective)

IMPLANT

How to use it:

- Placed by health care provider
- Lasts up to 3 years

Chances of getting pregnant: Less than 1 out of 100 women*

IUD

How to use it:

- Placed by health care provider
- Copper IUD lasts up to 10 years
- Progestin IUD lasts 3–5 years

Chances of getting pregnant: Less than 1 out of 100 women*

PILL

How to use it:

• Take at the same time each day

Chances of getting pregnant: 9 out of 100 women*

MALE CONDOM

How to use it:

• Use correctly every time during sex

Chances of getting pregnant: 18 out of 100 women*

Condoms should always be used along with the preferred birth control to protect against sexually transmitted diseases.

*Number of pregnancies per 100 women using the method within first year of typical use.

SOURCE: Trussell J., Contraception, May 2011 Last Updated Apr. 7, 2015