

# SPECIALTY GUIDELINE MANAGEMENT

## PROLIA (denosumab)

### POLICY

#### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### A. FDA-Approved Indications

1. Treatment of postmenopausal women with osteoporosis at high risk for fracture
2. Treatment to increase bone mass in men with osteoporosis at high risk for fracture
3. Treatment of men and women with glucocorticoid-induced osteoporosis at high risk for fracture
4. Treatment to increase bone mass in men at high risk for fracture receiving androgen deprivation therapy (ADT) for non-metastatic prostate cancer
5. Treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer

##### B. Compendial Uses

1. Prevention or treatment of osteoporosis during androgen deprivation therapy for prostate cancer in patients with high fracture risk
2. Consider in postmenopausal patients receiving adjuvant endocrine therapy along with calcium and vitamin D supplementation to maintain or improve bone mineral density and reduce risk of fractures

All other indications are considered experimental/investigational and not medically necessary.

#### II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review: Supporting chart notes or medical record indicating a history of fractures, T-score, and FRAX fracture probability as applicable to Sections III.A, III.B, and III.C.

#### III. CRITERIA FOR INITIAL APPROVAL

##### A. **Postmenopausal osteoporosis**

Authorization of 12 months may be granted to postmenopausal members with osteoporosis when ANY of the following criteria are met:

1. Member has a history of fragility fractures
2. Member has a pre-treatment T-score less than or equal to -2.5 OR member has osteopenia (i.e., pre-treatment T-score greater than -2.5 and less than -1) with a high pre-treatment FRAX fracture probability (See Appendix B) and meets ANY of the following criteria:

- a. Member has indicators of very high fracture risk (e.g., advanced age, frailty, glucocorticoid use, very low T-scores [less than or equal to -3], or increased fall risk)
- b. Member has failed prior treatment with or is intolerant to previous injectable osteoporosis therapy (e.g., zoledronic acid [Reclast], teriparatide [Forteo, Bonsity], abaloparatide [Tymlos])
- c. Member has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate (See Appendix A)

**B. Osteoporosis in men**

Authorization of 12 months may be granted to male members with osteoporosis when ANY of the following criteria are met:

1. Member has a history of an osteoporotic vertebral or hip fracture<sup>10</sup>
2. Member meets BOTH of the following criteria:
  - a. Member has a pre-treatment T-score less than or equal to -2.5 OR member has osteopenia (i.e., pre-treatment T-score greater than -2.5 and less than -1) with a high pre-treatment FRAX fracture probability (See Appendix B)<sup>10</sup>
  - b. Member has had an oral OR injectable bisphosphonate trial of at least 1-year duration OR there is a clinical reason to avoid treatment with a bisphosphonate (See Appendix A)

**C. Glucocorticoid-induced osteoporosis**

Authorization of 12 months may be granted to members with glucocorticoid-induced osteoporosis when ALL of the following criteria are met:

1. Member is currently receiving or will be initiating glucocorticoid therapy at an equivalent prednisone dose of  $\geq 2.5$  mg/day for  $\geq 3$  months.
2. Member has had an oral OR injectable bisphosphonate trial of at least 1-year duration OR there is a clinical reason to avoid treatment with a bisphosphonate (See Appendix A)
3. Member meets ANY of the following criteria:
  - a. Member has a history of a fragility fracture
  - b. Member has a pre-treatment T-score less than or equal to -2.5
  - c. Member has osteopenia (i.e., pre-treatment T-score greater than -2.5 and less than -1) with a high pre-treatment FRAX fracture probability (See Appendix B)

**D. Breast cancer**

Authorization of 12 months may be granted to members who are receiving adjuvant endocrine therapy for breast cancer.

**E. Prostate cancer**

Authorization of 12 months may be granted to members who are receiving androgen deprivation therapy for prostate cancer.

**IV. CONTINUATION OF THERAPY**

Authorization of 12 months may be granted for all members (including new members) who are currently receiving the requested medication through a previously authorized pharmacy or medical benefit, who meet one of the following:

1. Member has experienced clinical benefit as evidenced by a bone mass measurement showing an improvement or stabilization in T-score compared with the previous bone mass measurement and member has not experienced any adverse effects.

- Member has received less than 24 months of therapy and has experienced clinical benefit as evidenced by no adverse events during therapy (i.e., no clinically significant adverse reaction, no new fracture seen on radiography).

## V. APPENDIX

### Appendix A. Clinical reasons to avoid oral bisphosphonate therapy

- Presence of anatomic or functional esophageal abnormalities that might delay transit of the tablet (e.g. achalasia, stricture, or dysmotility Active upper gastrointestinal problem (e.g., dysphagia, gastritis, duodenitis, erosive esophagitis, ulcers)
- Presence of documented or potential gastrointestinal malabsorption (e.g. gastric bypass procedures, celiac disease, Crohn's disease, infiltrative disorders, etc.)
- Inability to stand or sit upright for at least 30 to 60 minutes
- Inability to take at least 30 to 60 minutes before first food, drink, or medication of the day
- Renal insufficiency (creatinine clearance <35 mL/min)
- History of intolerance to an oral bisphosphonate

### Appendix B. WHO Fracture Risk Assessment Tool

- High FRAX fracture probability: 10 year major osteoporotic fracture risk  $\geq$  20% or hip fracture risk  $\geq$  3%.
- 10-year probability; calculation tool available at: <https://www.sheffield.ac.uk/FRAX/>
- The estimated risk score generated with FRAX should be multiplied by 1.15 for major osteoporotic fracture (including fractures of the spine (clinical), hip, wrist, or humerus) and 1.2 for hip fracture if glucocorticoid treatment is greater than 7.5 mg (prednisone equivalent) per day.

## VI. REFERENCES

1. Prolia [package insert]. Thousand Oaks, CA: Amgen Inc.; March 2020.
2. The NCCN Drugs & Biologics Compendium™ © 2020 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed October 13, 2020.
3. Bisphosphonates. *Drug Facts and Comparisons*. Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; March 21, 2019. Accessed October 13, 2020.
4. Cosman F, de Beur SJ, LeBoff MS, et al. National Osteoporosis Foundation. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2014;25(10): 2359-2381.
5. Jeremiah MP, Unwin BK, Greenwald MH, et al. Diagnosis and management of osteoporosis. *Am Fam Physician*. 2015;92(4):261-268.
6. Camacho PM, Petak SM, Binkley N, et al. American Association of Clinical Endocrinologists and American College of Endocrinology Clinical Practice Guidelines for the Diagnosis and Treatment of Postmenopausal Osteoporosis 2020. *Endocr Pract*. 2020;26 (Suppl 1):1-46.
7. ACOG Practice Bulletin Number 129: Osteoporosis. *Obstet Gynecol*. 2012;120(3):718-734.
8. National Institute for Health and Care Excellence. Osteoporosis Overview. Last updated February 2018. Available at: <http://pathways.nice.org.uk/pathways/osteoporosis>. Accessed October 3, 2019.
9. Treatment to prevent osteoporotic fractures: an update. Department of Health and Human Services, Agency for Healthcare Research and Quality. 2012; Publication No. 12-EHC023-EF. Available at [www.effectivehealthcare.ahrq.gov/lbd.cfm](http://www.effectivehealthcare.ahrq.gov/lbd.cfm).
10. Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men: an Endocrine Society clinical practice guideline. *J Clin Endocr Metab*. 2012;97(6):1802-1822.
11. Gralow JR, Biermann S, Farooki A, et al. NCCN Task Force Report: Bone Health in Cancer Care. *JNCCN*. 2013; 11(Suppl 3):S1-50.

12. FRAX® WHO fracture risk assessment tool. © World Health Organization Collaborating Centre for Metabolic Bone Diseases: University of Sheffield, UK. Available at: <https://www.sheffield.ac.uk/FRAX/>. Accessed October 13, 2020.
13. Fink HA, Gordon G, Buckley L, et al. 2017 American College of Rheumatology Guidelines for the Prevention and Treatment of Glucocorticoid-Induced Osteoporosis. *Arthritis Care Res.* 2017;69:1521-1537.
14. Ensrud KE, Crandall CJ. Osteoporosis. *Ann Intern Med* 2017;167(03):ITC17–ITC32.
15. Eastell R, Rosen CJ, Black DM, et al. Pharmacological Management of Osteoporosis in Postmenopausal Women: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2019;104:1595-1622.
16. Carey John. What is a 'failure' of bisphosphonate therapy for osteoporosis? *Cleveland Clinic Journal of Medicine* Nov 2005, 72 (11) 1033-1039.