



# Preoperative Bathing Instructions

Before surgery, you can play an important role in your health. Because skin is not sterile, you can reduce the number of germs on your skin by carefully washing before surgery. Please follow these instructions.

**IMPORTANT:** You will need to shower with a special soap called chlorhexidine gluconate (CHG). A common brand name for this soap is Hibiclens, but any brand is acceptable. You will only need a small bottle, about 8 ounces.

## Bathing Instructions

- You will need to take **three** CHG showers. The first shower is 2 days before surgery, then the night before surgery, and the morning of surgery.
- With each shower, wash your hair as usual with your normal shampoo. Rinse your hair and body thoroughly afterward to remove the residue.
- Do not shave the area of your body where your surgery will be performed.
- Turn the water off before applying the CHG soap to prevent rinsing it off too soon. Apply CHG to your entire body from the jaw down. Do not get the soap in your eyes, ears, nose, mouth or genital area. Wash thoroughly for five minutes, paying special attention to the area where your surgery will be performed. Do not scrub your skin too hard. Do not wash with your regular soap after using CHG. Turn the water back on and rinse your body well.
- Pat yourself dry with a clean, soft towel after each shower. Then put on clean clothes or pajamas and sleep on freshly laundered bed linens. Do not apply any lotions, roll on deodorants, perfumes or powders after using CHG the day of surgery.
- If allergic to Chlorhexidine, use antibacterial soap (such as Dial) for your shower.